arXiv:math/0510268v1 [math.DG] 13 Oct 2005

Prequantization of the moduli space of flat connections over a four-manifold

Tosiaki Kori[*]
Department of Mathematics
School of Science and Engineering
Waseda University
3-4-1 Okubo, Shinjuku-ku Tokyo, Japan.
e-mail: kori@waseda.jp

Abstract

We introduce a symplectic structure on the space of connections in a G-principal bundle over a four-manifold and the Hamiltonian action on it of the group of gauge transformations which are trivial on the boundary. The symplectic reduction becomes the moduli space of flat connections over the manifold. On the moduli space of flat connections we shall construct a hermitian line bundle with connection whose curvature is given by the symplectic form. This is the Chern-Simons prequantum line bundle. The group of gauge transformations on the boundary of the base manifold acts on the moduli space of flat connections by an infinitesimally symplectic way. This action is lifted to the prequantum line bundle by its abelian extension.

MSC: 57R; 58E; 81E.
Subj. Class: Global analysis, Quantum field theory.
Keywords Chern-Simons prequantization. Wess-Zumino-Witten actions. Four-manifolds.

0 Introduction

Atiyah and Bott [2] showed that the moduli space of flat connections on the trivial $SU(2)$ bundle over a surface Σ is a compact, finite-dimensional

---

[*] Research supported by Promotion for Sciences of the Ministry of Education and Science in Japan (no. 16540084)

symplectic space. Ramadas, Singer and Weitsman [13] described the Chern-Simons prequantization of this moduli space. They showed heuristically how the path-integral quantization of the Chern-Simons action yields holomorphic sections of the prequantum line-bundle when Σ is endowed with a complex structure. On the other hand, Donaldson proved that if a two-manifold has the boundary then the moduli space of flat connections is a smooth infinite-dimensional symplectic manifold, and it has a Hamiltonian group action of the gauge transformations on the boundary [6].

In general the moduli space of the $Lie(G)$-valued flat connections on a a connected compact manifold M without boundary corresponds bijectively to the conjugate classes of the G-representations of the fundamental group $\pi_1(M)$. In this paper we study the Chern-Simons prequantization of the space of flat connections on a four manifold M generally with non-empty boundary.

Let $\mathcal{A}(M)$ be the space of irreducible connections on M and let $\mathcal{G}_0(M)$ be the group of gauge transformations on M that reduce to the identity transformation when restricted to the boundary ∂M. We shall prove in section 1 that $\mathcal{A}(M)$ carries a symplectic structure. The symplectic form is given by

$$\begin{aligned} \omega_A(a,b) &= \frac{1}{8\pi^3}\int_M Tr[\,(ab-ba)F_A\,] - \frac{1}{24\pi^3}\int_{\partial M} Tr[\,(ab-ba)\,A\,] \\ &\qquad + \int_M Tr[\,G_A(*[a,*b])\,]dV\,, \end{aligned}$$

for a, $b \in T_A\mathcal{A}$. Where $G_A = (\Delta_A)^{-1}$ is the Green operator of the Dirichlet problem. The action of $\mathcal{G}_0(M)$ becomes a Hamiltonian action with the moment map given by the square of curvature F_A^2. Hence the 0-level set of the moment map is the space of flat connections $\mathcal{A}^\flat(M)$. Let $\mathcal{M}^\flat(\partial M)$ be the symplectic reduction, that is, the orbit space of $\mathcal{A}^\flat(M)$ by $\mathcal{G}_0(M)$. We prefer to adopt the notation $\mathcal{M}^\flat(\partial M)$ than to write $\mathcal{M}^\flat(M)$ because $\mathcal{A}^\flat(M)/\mathcal{G}_0(M)$ is near to $\mathcal{A}^\flat(\partial M)$, the space of flat connections over ∂M.

In section 2 we shall investigate the Chern-Simons prequantization of four manifolds. By a *prequantization* over M we mean a hermitian line bundle $\mathcal{L}(\partial M)$ with connection over $\mathcal{M}^\flat(\partial M)$ whose curvature is given by the symplectic form of $\mathcal{M}^\flat(\partial M)$, [3, 13]. When M has no boundary $\mathcal{L}(\partial M)$ becomes a vector space and the *Chern-Simons functional* over M is defined as a vector of this space. When M has the boundary the Chern-Simons functional over M is defined as a section of the pull back line bundle of $\mathcal{L}(\partial M)$ by the quotient map $p : \mathcal{A}^\flat(M) \longrightarrow \mathcal{M}^\flat(\partial M)$. The Chern-Simons functional gives a non-vanishing horizontal section over of the pullback line bundle $p^*\mathcal{L}(\partial M)$.

The set of line bundles $\mathcal{L}(\partial M)$ and the associated Chern-Simons functionals CS_M form a category that represents the operations of union and contraction on the cobordism classes of base manifolds M, that is, they respect the axioms of Topological Field Theory [1, 8]. We describe our *Chern-Simons prequantization* (CSQ) as a functor CS from the category of some classes of four-manifolds with boundary to the category of complex line bundles:

$$(\,\Gamma = \partial M,\, M\,) \Longrightarrow (\,\mathcal{L}(\Gamma),\, CS_M\,).$$

As for the axioms of CSQ, we require other than the axioms of TFT two axioms that are characteristic to CSQ. The first says that $\pi : \mathcal{L}(\Gamma) \longrightarrow \mathcal{M}^\flat(\Gamma)$ is a hermitian line bundle with connection whose curvature is equal to the symplectic form on $\mathcal{M}^\flat(\Gamma)$, and the second says that $\exp 2\pi i\, \mathrm{CS}_M$ is a non-vanishing horizontal section of $p^*\mathcal{L}(\Gamma)$.

We shall construct the Chern-Simons prequantization explicitly for $M = S^4$ and for $M = D^4$, the four-dimensional hemisphere. Then we give by a functorial method the prequantization over a conformally flat four-manifold with boundary.

The group of pointed gauge transformations $\mathcal{G}(M)$ acts on $\mathcal{A}^\flat(M)$ by infinitesimal symplectic automorphisms. So is the action of $\mathcal{G}(\partial M)$ on $\mathcal{M}^\flat(\partial M)$. We shall discuss in section 3 the lift of this action to the prequantum line bundle $\mathcal{L}(\partial M)$. We prove that the abelian extension of $\mathcal{G}(\partial M)$ introduced by Mickelsson [10] acts on $\mathcal{L}(\partial M)$. As was discussed in [8] the geometric description of this abelian extension is given by the four-dimensional Wess-Zumino-Witten model.

1 Symplectic structure on the space of connections

1.1 Differential calculation on $\mathcal{A}$

Let M be an oriented Riemannian four-manifold with boundary ∂M. Let $G = SU(N)$. The inner product on G is given by $<\xi, \eta>= -Tr(\xi, \eta)$, $\xi,\, \eta \in Lie(G) = su(N)$. . With this inner product the dual of $Lie(G)$ is identified with $Lie(G)$ itself. Let $\pi : P \longrightarrow M$ be a principal G-bundle over M which is given by a system of transition functions in the Sobolev space L^2_s for $s > 2$. We write $\mathcal{A} = \mathcal{A}(M)$ for the space of *irreducible* L^2_{s-1} connections, which differ from a smooth connection by an L^2_{s-1} 1-form on P with values in $Lie(G)$, hence the tangent space of $\mathcal{A}$ at $A \in \mathcal{A}$ is $T_A\mathcal{A} = \Omega^1_{s-1}(M, adP)$. Similarly we introduce the space of connections $\mathcal{A}(\partial M)$. It is the $L^2_{s-\frac{1}{2}}$

connections that differ from a smooth connection on ∂M by a $L^2_{s-\frac{1}{2}}$ 1-form on ∂M. The boundary restriction map $r : \mathcal{A}(M) \longrightarrow \mathcal{A}(\partial M)$ is surjective.

The curvature of $A \in \mathcal{A}$ is

$$F_A = dA + \frac{1}{2}[A \wedge A] \in \Omega^2_{s-2}(M, adP).$$

$\Omega^3_{s-2}(M, adP)$, being the dual of $\Omega^1_{s-1}(M, adP) = T_A\mathcal{A}$, is identified with the space of 1-forms on $\mathcal{A}$.

Here are some differential calculations on $\mathcal{A}$ that we shall cite from [4, 7, 14].

The derivation of a smooth function $G = G(A)$ on $\mathcal{A}$ is defined by the functional variation of A:

$$(\delta_A G)a = \lim_{t \longrightarrow 0} \frac{G(A + ta) - G(A)}{t}, \qquad \text{for } a \in T_A\mathcal{A}. \tag{1.1}$$

We have, for example,

$$(\delta_A A)a = a, \qquad (\delta_A F_A)a = d_A a.$$

The second follows from the formula

$$F_{A+a} = F_A + d_A a + a \wedge a.$$

Similarly the derivation of a vector field on $\mathcal{A}$ or a 1-form on $\mathcal{A}$ is defined as that of a smooth function of $A \in \mathcal{A}$ valued in $\Omega^1(M, adP)$, respectively, in $\Omega^3(M, adP)$. We have, for a vector field $\mathbf{b}$ and a 1-form β,

$$(\delta_A < \beta, \mathbf{b} >)a =< \beta, (\delta_A \mathbf{b})a > + < (\delta_A \beta)a, \mathbf{b} > . \tag{1.2}$$

where $< \beta, \mathbf{b} >$ is the exterior product of $Lie(G)$ valued forms on M.

The Lie bracket for vector fields on $\mathcal{A}$ is seen to have the expression

$$[\mathbf{a}, \mathbf{b}] = (\delta_A \mathbf{b})\mathbf{a} - (\delta_A \mathbf{a})\mathbf{b}. \tag{1.3}$$

Let $\widetilde{d}$ be the exterior derivative on $\mathcal{A}$. For a function on $\mathcal{A}$, $(\widetilde{d}G)_A a = (\delta_A G)a$. From (1.2) and (1.3) we have the following formula for the exterior derivative of a 1-form on $\mathcal{A}$:

$$\begin{aligned} (\widetilde{d}\theta)_A(\mathbf{a}, \mathbf{b}) &= (\delta_A < \theta, \mathbf{b} >)\mathbf{a} - (\delta_A < \theta, \mathbf{a} >)\mathbf{b} - < \theta, [\mathbf{a}, \mathbf{b}] > \\ &= < (\delta_A \theta)\mathbf{a}, \mathbf{b} > - < (\delta_A \theta)\mathbf{b}, \mathbf{a} > . \end{aligned} \tag{1.4}$$

Likewise, if ω is a 2-form on $\mathcal{A}$, then

$$(\widetilde{d}\omega)_A(\mathbf{a}, \mathbf{b}, \mathbf{c}) = (\delta_A \omega(\mathbf{b}, \mathbf{c}))\mathbf{a} + (\delta_A \omega(\mathbf{c}, \mathbf{a}))\mathbf{b} + (\delta_A \omega(\mathbf{c}, \mathbf{a}))\mathbf{b}. \tag{1.5}$$

1.2 Gauge transformations

We write $\mathcal{G}' = \mathcal{G}'(M)$ for the group of L^2_s gauge transformations:

$$\mathcal{G}'(M) = \Omega^0_s(M, AdP). \tag{1.6}$$

$\mathcal{G}'$ acts on $\mathcal{A}$ by

$$g \cdot A = g^{-1}dg + g^{-1}Ag = A + g^{-1}d_A g. \tag{1.7}$$

By Sobolev lemma one sees that $\mathcal{G}'$ is a Banach Lie group and the action is a smooth map of Banach manifolds.

The group of $L^2_{s-1/2}$ gauge transformations on the boundary ∂M is denoted by $\mathcal{G}'(\partial M)$. We have the restriction map to the boundary:

$$r : \mathcal{G}'(M) \longrightarrow \mathcal{G}'(\partial M).$$

Let $\mathcal{G}_0 = \mathcal{G}_0(M)$ be the kernel of the restriction map. It is the group of gauge transformations that are identity on the boundary. $\mathcal{G}_0$ acts freely on $\mathcal{A}$ and $\mathcal{A}/\mathcal{G}_0$ is therefore a smooth infinite dimensional manifold, while the action of $\mathcal{G}'$ is not free. In the following we shall choose $p_0 \in M$ fixed point on the boundary ∂M and deal with the group of gauge transformations leaving p_0 fixed:

$$\mathcal{G} = \mathcal{G}(M) = \{g \in \mathcal{G}'(M);\ g(p_0) = 1\}.$$

If $\partial M = \phi$, p_0 is any point of M . $\mathcal{G}$ act freely on $\mathcal{A}$. Let $\mathcal{A}/\mathcal{G}$ be the orbit space of this action. It is a smooth infinite dimensional manifold. We have $Lie(\mathcal{G}) = \Omega^0(M, adP)$.

Correspondingly we have the group $\mathcal{G}(\partial M) = \{g \in \mathcal{G}'(\partial M);\ g(p_0) = 1\}$, and the restriction map $r : \mathcal{G}(M) \longrightarrow \mathcal{G}(\partial M)$ with the kernel $\mathcal{G}_0(M)$. We have

$$Lie(\mathcal{G}_0) = \{\xi \in Lie(\mathcal{G});\ \xi|\partial M = 0\}. \tag{1.8}$$

The derivative of the action of $\mathcal{G}$ at $A \in \mathcal{A}$ is

$$d_A = d + [A\wedge \quad] : \Omega^0(M, adP) \longrightarrow \Omega^1(M, adP). \tag{1.9}$$

Thus the fundamental vector field on $\mathcal{A}$ corresponding to $\xi \in Lie(\mathcal{G})$ is given by

$$\xi_{\mathcal{A}}(A) = \frac{d}{dt}|_{t=0}(\exp t\xi) \cdot A = d_A\xi.$$

So the tangent space to the orbit at $A \in \mathcal{A}$ is

$$T_A(\mathcal{G} \cdot A) = \{d_A\xi;\ \xi \in \Omega^0(M, adP)\} \tag{1.10}$$

We have two orbit spaces;

$$\mathcal{B}(M) = \mathcal{A}/\mathcal{G}_0, \qquad \mathcal{C}(M) = \mathcal{A}/\mathcal{G}. \tag{1.11}$$

1.3 Canonical connections on the $\mathcal{G}_0$-orbit space and $\mathcal{G}$-orbit space.

Let $\mathcal{A} = \mathcal{A}(M)$, $\mathcal{G} = \mathcal{G}(M)$, $\mathcal{B} = \mathcal{B}(M)$ and $\mathcal{C} = \mathcal{C}(M)$ be as stated above. We shall investigate the horizontal subspaces of the fibrations $\mathcal{A} \longrightarrow \mathcal{B}$ and that of $\mathcal{A} \longrightarrow \mathcal{C}$.

We have the Stokes formula

$$\int_{\partial M} f * u = \int_M d_A f \wedge *u - \int_M f * d_A^* u,$$

for $f \in \Omega^0(M, adP)$, $u \in \Omega^1(M, adP)$. Hence we see that $d_A\, Lie(\mathcal{G}_0)$ is orthogonal to the space $\ker d_A^* = \{a \in \Omega^1(M, adP);\, d_A^* a = 0\}$, and .that $d_A\, Lie(\mathcal{G})$ is orthogonal to $\{a \in \ker d_A^*,\ (*a)|\partial M = 0\}$

Let Δ_A be the covariant Laplacian defined as the closed extension of $d_A^* d_A$ with the domain of definition $\mathcal{D}_{\Delta_A} = \{u \in \Omega^0_s(M, adP);\, u|\partial M = 0\}$. Since $A \in \mathcal{A}$ is irreducible $\Delta_A : \mathcal{D}_{\Delta_A} \longrightarrow \Omega^0_{s-2}(M, adP)$ is an isomorphism. Let $G_A = (\Delta_A)^{-1}$ be the Green operator of the Dirichlet problem :

$$\begin{cases} \Delta_A u &= f \\ u|\partial M &= 0 \end{cases}$$

Proposition 1.1. *Let $A \in \mathcal{A}$.*

1. *We have the following orthogonal decomposition:*

   $$T_A\mathcal{A} = \{d_A\xi;\, \xi \in Lie(\mathcal{G}_0)\,\} \,\oplus\, H_A^0, \tag{1.12}$$

   where

   $$H_A^0 = \{a \in \Omega^1(M, adP);\, d_A^* a = 0\}.$$

2. *The $\mathcal{G}_0$-principal bundle $\pi : \mathcal{A} \longrightarrow \mathcal{B}$ has a natural connection defined by the horizontal subspace H_A^0, which is given by the connection form $\gamma_A^0 = G_A d_A^*$.*

3. *The curvature form $\mathcal{F}^0$ of the connection form γ^0 is given by*

   $$\mathcal{F}_A^0(a, b) = G_A(*\,[a,\, *b]) \qquad \text{for } a, b \in H_A^0.$$

When M is compact and $\partial M = \phi$, the proposition is well known [5, 14]., and ours are proved by the same argument.

Corollary 1.2.

$$\mathcal{F}^0_A(a, d_A\xi) = 0 \qquad \textit{for } \xi \in Lie(\mathcal{G}_0). \tag{1.13}$$

Now we proceed to the fibration $\mathcal{A} \longrightarrow \mathcal{C} = \mathcal{A}/\mathcal{G}$.

Fow a 1-form v, let $g = K_A v$ denote the solution of the following boundary value problem:

$$\left\{ \begin{array}{ccc} \Delta_A g & = & 0 \\ *d_A g|\partial M & = & *v|\partial M. \end{array} \right.$$

Proposition 1.3. *Let $A \in \mathcal{A}$.*

1. *We have the orthogonal decomposition:*

$$T_A\mathcal{A} = \{d_A\xi;\ \xi \in Lie(\mathcal{G})\} \ \oplus\ H_A, \tag{1.14}$$

where

$$H_A = \{a \in \Omega^1(M, adP);\ d^*_A a = 0,\ \textit{and}\ *a|\partial M = 0\}$$

2. *The $\mathcal{G}$-principal bundle $\pi : \mathcal{A} \longrightarrow \mathcal{C}$ has a natural connection defined by the horizontal subspace H_A.*

Proof

Let $a \in \Omega^1(M, adP)$ and $a = d_A\xi + b$ be the decomposition of (1.11), then $\xi = G_A d^*_A a$ and $b \in \Omega^1(M, adP)$, $d^*_A b = 0$. Put $\eta = K_A b$. Then we have the orthogonal decomposition

$$a = d_A(\xi + \eta) + c,$$

with $c \in H_A$ and $\xi + \eta \in Lie(\mathcal{G})$. If we write

$$\gamma_A = \gamma^0_A + K_A(I - d_A\gamma^0_A), \tag{1.15}$$

where I is the identity transformation on $T_A\mathcal{A}$, then γ_A is a $Lie(\mathcal{G})-$valued 1-form which vanishes on H_A and $\gamma_A d_A\xi = \xi$, that is, γ_A is the connection form. □

Let $g = N_A f$ be the solution of Neuman problem:

$$\left\{ \begin{array}{ccc} \Delta^n_A g & = & f \\ *d_A g|\partial M & = & 0 \qquad \text{on } \partial M. \end{array} \right.$$

Where Δ^n_A is the closed extension of $d^*_A d_A$ with the domain of definition $\mathcal{D}_{\Delta^n_A} = \{u \in \Omega^0_s(M, adP);\ *d_A u|\partial M = 0\}$.

Corollary 1.4. *The curvature form is given by*

$$\mathcal{F}_A(a,b) = N_A(*[a,\, *b]) \qquad \textit{for } a,b \in H_A. \tag{1.16}$$

1.4 Symplectic structure on $\mathcal{A}$

For each $A \in \mathcal{A}(M)$ we define a sqew-symmetric bilinear form on $T_A\mathcal{A}$ by:

$$\begin{aligned}
\omega_A(a,b) &= \omega_A^0(a,b) + \omega_A'(a,b) + \phi_A(a,b), && (1.17)\\
\omega_A^0(a,b) &= \frac{1}{8\pi^3}\int_M Tr[\,(a\wedge b - b\wedge a)\wedge F_A\,], && (1.18)\\
\omega_A'(a,b) &= -\frac{1}{24\pi^3}\int_{\partial M} Tr[\,(a\wedge b - b\wedge a)\wedge A\,], && (1.19)\\
\phi_A(a,b) &= \int_M Tr[\,\mathcal{F}_A^0(a,b)\,]dx, && (1.20)
\end{aligned}$$

for $a,b \in T_A\mathcal{A}$. Here $\mathcal{F}^0$ is the curvature form of Proposition 1.1 and dx is the volume form on M.

Evidently ω is non-degenerate.

Theorem 1.5. *$(\mathcal{A}(M),\omega)$ is a symplectic space.*

Proof

In the following we shall abbreviate ab for the exterior product $a \wedge b$. From the Bianchi's identity we have $\widetilde{d}\mathcal{F}^0 = 0$, hence $\widetilde{d}\phi = 0$.

Differentiating the 2-form ω^0, we have

$$(\widetilde{d}\omega^0)_A(a,b,c) = \delta_A(\omega^0(a,b))(c) + \delta_A(\omega^0(b,c))(a) + \delta_A(\omega^0(c,a))(b),$$

for $a,b,c \in T_A\mathcal{A}$. From the definition we have

$$\delta_A(\omega^0(a,b))(c) = \frac{1}{8\pi^3}\int_M Tr[(ab-ba)d_Ac],$$

hence

$$(\widetilde{d}\omega^0)_A(a,b,c) = \frac{1}{8\pi^3}\int_M Tr\left[(ab-ba)d_Ac + (bc-cb)d_Aa + (ca-ac)d_Ab\right].$$

Since

$$d(Tr(ab-ba)c) = Tr\left[(ab-ba)d_Ac + (bc-cb)d_Aa + (ca-ac)d_Ab\right],$$

we have

$$(\widetilde{d}\omega^0)_A(a,b,c) = \frac{1}{8\pi^3}\int_M d(Tr(ab-ba)c) = \frac{1}{8\pi^3}\int_{\partial M} Tr[(ab-ba)c].$$

On the other hand we have

$$(\widetilde{d}\omega')_A(a,b,c) = 3\delta_A(\omega'(a,b))(c) = -\frac{1}{8\pi^3}\int_{\partial M} Tr[(ab-ba)c].$$

Therefore $\widetilde{d}\omega = 0$.

Thus $(\mathcal{A},\omega)$ is a symplectic manifold. □

Corollary 1.6. *$\mathcal{G}_0$ acts symplectically on $\mathcal{A}$.*

Note that the action of $\mathcal{G}$ is not even infinitesimally symplectic.

Proposition 1.7. *The canonical 1-form θ that corresponds to the symplectic form ω ;*

$$(\widetilde{d}\theta)_A = \omega_A,$$

is given by

$$\theta_A(a) = -\frac{1}{24\pi^3}\int_M Tr[(AF + FA - \frac{1}{2}A^3)\,a + \int_M Tr[\gamma_A\,a]dx. \qquad (1.21)$$

Remark 1.1. For M without boundary the pre-symplectic form ω^0 and the Hamiltonian action of $\mathcal{G} = \mathcal{G}_0$ in the next theorem were introduced by Bao and Nair [4]. More generally they gave the pre-symplectic form on n-dimensional manifolds.

1.5 Flat connections

The space of flat connections are denoted by

$$\mathcal{A}^\flat(M) = \{A \in \mathcal{A}(M); F_A = 0\},$$

which we shall often abbreviate to $\mathcal{A}^\flat$. The tangent space of $\mathcal{A}^\flat$ is given by

$$T_A\mathcal{A}^\flat = \{a \in \Omega^1(M, adP); d_A a = 0\}.$$

Here the Sobolev indeces are suppressed.

Theorem 1.8. *The action of $\mathcal{G}_0$ on $\mathcal{A}$ is a Hamiltonian action and the corresponding moment map is given by*

$$\Phi \quad : \quad \mathcal{A} \longrightarrow (Lie\,\mathcal{G}_0)^* = \Omega^4(M, adP) : \qquad A \longrightarrow F_A^2.$$

$$\langle \Phi(A), \xi\rangle \quad = \quad \Phi^\xi(A) = \frac{1}{8\pi^3}\int_M Tr(F_A^2\xi) \qquad (1.22)$$

Proof

We have

$$(\delta\Phi^{\xi})_A a = \frac{1}{8\pi^3}\int_M Tr[(d_A a \wedge F_A + F_A \wedge d_A a)\xi],$$

and

$$\begin{aligned}(\delta\Phi^{\xi})_A a \quad - \quad \omega^0_A(a, d_A\xi) &= \frac{1}{8\pi^3}\int_M Tr[(d_A a \wedge F_A + F_A \wedge d_A a)\xi - (F_A a + aF_A)d_A\xi] \\ &= \frac{1}{8\pi^3}\int_M dTr[(aF_A + F_A a)\xi] = \frac{1}{8\pi^3}\int_{\partial M} Tr[(aF_A + F_A a)\xi] = 0,\end{aligned}$$

since $\xi = 0$ on ∂M. On the other hand we have

$$\begin{aligned}dTr[(Aa - aA)\xi] \quad &= \quad Tr[(F_A a + aF_A - Ad_A a - d_A aA + A^2 a + aA^2)\xi] + Tr[(Aa - aA)d_A\xi] \\ &= Tr[(ad_A\xi - d_A\xi a)A]\end{aligned}$$

on ∂M. Hence

$$\omega'(a, d_A\xi) = -\frac{1}{24\pi^3}\int_{\partial M} dTr[(Aa - aA)\xi] = 0.$$

We have also $\phi(a, d_A\xi) = 0$ from Corollary 1.2. Therefore

$$(\delta\Phi^{\xi})_A a = \omega_A(a, d_A\xi).$$

The moment map for the action of $\mathcal{G}_0$ on $\mathcal{A}$ is

$$\Phi : \ A \longrightarrow F_A^2$$

□

The symplectic quotient of $\mathcal{A}$ by $\mathcal{G}_0$ is

$$\mathcal{M}^{\flat} = \Phi^{-1}(0)/\mathcal{G}_0 = \mathcal{A}^{\flat}(M)/\mathcal{G}_0.$$

Remark 1.2. $\mathcal{M}^{\flat}$ is the conjugacy classes of G-representations of $\pi_1(M, \partial M)$.

Theorem 1.9. *$\mathcal{M}^{\flat}$ is a smooth symplectic manifold. The symplectic form on $\mathcal{M}^{\flat}$ is given by*

$$\omega^{\flat}_{[A]}([a], [b]) = \omega'_A(a, b) \tag{1.23}$$

for $[A] \in \mathcal{M}^{\flat}$ and $[a]$, $[b] \in T_{[A]}\mathcal{M}^{\flat}$.

This is the symplectic reduction theorem of Marsden-Weinstein [9] . Let $(\mathcal{G}_0)_A$ be the isotropy group of A, then the image by $(\delta\Phi)_A$ of $T_A\mathcal{A}$ is the annihilator of

$$Lie((\mathcal{G}_0)_A) = \{\xi \in \Omega^0(M, adP);\ d_A\xi = 0,\ \xi|\partial M = 0\} = 0,$$

since $A \in \mathcal{A}$ is irreducible. Hence $(\delta\Phi)_A$ is surjective. In particular $0 \in (Lie\,\mathcal{G}_0)^*$ is a regular value. Thus by the reduction theorem we see that $\mathcal{A}^\flat = \Phi^{-1}(0)$ is a co-isotropic submanifold of $\mathcal{A}$ and the orbit space $\mathcal{M}^\flat = \mathcal{A}^\flat/\mathcal{G}_0$ is a smooth manifold. The coordinate mappings are described by the implicit function theorem [7]. For $A \in \mathcal{A}^\flat$ there is a slice for the $\mathcal{G}_0$-action on $\mathcal{A}^\flat$ given by the Coulomb gauge condition:

$$V_A \subset \{A + a;\ F_{A+a} = 0,\ d_A^* a = 0\}.$$

V_A is a smooth Banach manifold homeomorphic to the following subset of the tangent space at A:

$$H_A \cap T_A\mathcal{A}^\flat = \{a \in \Omega^1(M, adP) :\ d_Aa = 0,\ d_A^*a = 0\}.$$

From the definition $\gamma_A^0 a = 0$ for $a \in H_A$. Thus there exists a symplectic structure on $\mathcal{M}^\flat$ given by

$$\omega^\flat_{[A]}([a],[b]) = \omega'_A(a,b). \tag{1.24}$$

Here $[A] \in \mathcal{M}^\flat$ denotes the $\mathcal{G}_0$-orbit of $A \in \mathcal{A}^\flat$, and for $[a] \in T_{[A]}\mathcal{M}^\flat$ we take the representative tangent vector to the slice; $a \in H_A \cap T_A\mathcal{A}^\flat$. $\omega^\flat$ is well defined because we have $g \cdot A = A$ on ∂M for $g \in \mathcal{G}_0$, and $\omega'_A(a, d_A\xi) = 0$ for $\xi \in Lie\,\mathcal{G}_0$ and $a \in T_A\mathcal{A}^\flat$. □

Corollary 1.10.

$$T_{[A]}\mathcal{M}^\flat = \{a \in \Omega^1(M.adP);\ d_Aa = d_A^*a = 0, \} \tag{1.25}$$

In the following, when there is a doubt about which manifold is involved, we shall write $\mathcal{M}^\flat(\partial M)$ for the orbit space $\mathcal{M}^\flat = \mathcal{A}^\flat(M)/\mathcal{G}_0(M)$.

Example 1.

For $M = S^4$, $\partial S^4 = \emptyset$, we have $\mathcal{G}_0(S^4) = \mathcal{G}(S^4)$. The moduli space of flat connections is

$$\mathcal{M}^\flat(\emptyset) = \mathcal{A}^\flat(S^4)/\mathcal{G}(S^4).$$

Then $\mathcal{M}^\flat(\emptyset)$ is one-point.

In fact, let $p_0 \in S^4$ and let $A \in \mathcal{A}^\flat(S^4)$. Let $T^A_\gamma(x)$ denote the parallel transformation by A along the curve γ joining p_0 and x. We put $f_A(x) =$

$T^A_\gamma(x)1 \in G$. It is independent of the choice of curve γ joining p_0 and x. Then $f_A \in \mathcal{G}(S^4)$. By the definition $A = df_A \cdot f_A^{-1}$.

In general $\mathcal{M}^\flat(\emptyset)$ is one-point for a connected and simply connected compact manifold M without boundary.

Example 2.

For a disc $D = \{x \in \mathrm{R}^4; \, |x| \leq 1\}$ with boundary S^3, we have $\mathcal{M}^\flat(S^3) \simeq \Omega^3_0 G$. Where

$$\Omega^3 G = \{f \in Map(S^3, G); f(p_0) = 1\},$$

and $\Omega^3_0 G$ is its connected component of the identity. First we note that

$$\mathcal{G}(D) \simeq DG = \{f \in Map(D, G); \; f(p_0) = 1\},$$

and

$$\mathcal{G}_0(D) \simeq D_0 G = \{f \in DG; \; f|S^3 = 1\}.$$

Hence $\mathcal{G}/\mathcal{G}_0 \simeq \Omega^3_0 G$.

As before we put, for $A \in \mathcal{A}^\flat(D)$, $f_A(x) = T^A_\gamma(x)1$, $x \in D$. We have a well defined bijective map from $\mathcal{A}^\flat(D)$ to DG. In particular, $f_A = g$ for $A = dg\, g^{-1}$ with $g \in \mathcal{G}_0$. It holds also that $f_{g\cdot A}(x) = f_A(x)g(x)$ for $g \in \mathcal{G}$. Hence we have the isomorphism

$$\mathcal{M}^\flat(S^3) = \mathcal{A}^\flat/\mathcal{G}_0 \simeq DG/D_0G \simeq \Omega^3_0 G.$$

More generally we have $\mathcal{M}^\flat(\partial M) \simeq \mathcal{G}/\mathcal{G}_0$ when M is a manifold with boundary ∂M such that M is made into a simply connected compact manifold by capping a disc to each boundary component.

Now let $D' \subset S^4$ be the complementaly hemisphere of $D^4 \subset S^4$, the boundary $\partial D'$ is S^3 with the reversed orientation that we denote by $(S^3)' = \partial D'$. As above we have

$$\mathcal{M}^\flat((S^3)') = \mathcal{A}^\flat(D')/\mathcal{G}_0(D').$$

Lemma 1.11. *There is an isomorphism*

$$p : \mathcal{M}^\flat(S^3) \longrightarrow \mathcal{M}^\flat((S^3)'). \tag{1.26}$$

Proof

We define the map

$$p : \mathcal{A}^\flat(D) \longrightarrow \mathcal{M}^\flat((S^3)'). \tag{1.27}$$

by

$$p(A) = [A'],$$

where, for $A \in \mathcal{A}^\flat(D)$ we associated a $A' \in \mathcal{M}^\flat(D')$ such that $A|S^3 = A'|S^3$. The equivalence class $p(A) = [A']$ is well defined because, for another A'' with $A|S^3 = A''|S^3$, there is a $g' \in \mathcal{G}_0(D')$ such that $A'' = g' \cdot A'$. If $p(A) = p(B) = [A']$ then $A|S^3 = A'|S^3 = B|S^3$, so there is a $g \in \mathcal{G}_0(D)$ such that $B = g \cdot A$. Hence the kernel of p is $\mathcal{G}_0(D)$. Thus p induces an isomorphism:

$$p : \mathcal{M}^\flat(S^3) \longrightarrow \mathcal{M}^\flat((S^3)'). \tag{1.28}$$

□

The boundary restriction map $r : \mathcal{A}(M) \longrightarrow \mathcal{A}(\partial M)$ induces a map

$$\bar{r} : \mathcal{M}^\flat(\partial M) = \mathcal{A}^\flat(M)/\mathcal{G}_0 \longrightarrow \mathcal{A}^\flat(\partial M).$$

If A_1, $A_2 \in \mathcal{A}^\flat(M)$ have the same boundary restriction, then the corresponding holonomy groups coincide on the boundary, and defines a $g \in \mathcal{G}_0$ such that $g \cdot A_1 = A_2$. Therefore $\bar{r}$ is a injective map.

1.6 The action of $\mathcal{G}(M)$

By the action of the group of gauge transformations $\mathcal{G}$ on $\mathcal{A}^\flat$ we have the orbit space $\mathcal{N}^\flat = \mathcal{A}^\flat/\mathcal{G}$. Then we have a fibration $\mathcal{M}^\flat \longrightarrow \mathcal{N}^\flat$ with the fiber $\mathcal{G}(\partial M) = \mathcal{G}/\mathcal{G}_0$. We note the fact that any vector which is tangent to the $\mathcal{G}$-orbit through $A \in \mathcal{A}^\flat$ is in $T_A\mathcal{A}^\flat$. Propositions 1.3 yields the following Proposition.

Proposition 1.12. *Let $A \in \mathcal{M}^\flat$.*

1. *We have the following decomposition*

$$T_A\mathcal{M}^\flat \;=\; \{d_A\xi;\ \xi \in Lie(\mathcal{G}(\partial M))\} \oplus H^\flat_A, \tag{1.29}$$

where

$$H^\flat_A \;=\; \{a \in \Omega^1(M, adP);\ d_A a = d_A^* a = 0,\ \textit{and}\ *a|\partial M = 0\}.$$

2. *The $\mathcal{G}(\partial M)$-principal bundle $\pi : \mathcal{M}^\flat \longrightarrow \mathcal{N}^\flat$ has a natural connection defined by the horizontal space $H^\flat_A$. The connection form is given by $K_A(I - d_A\gamma^0_A)$.*

The action of $\mathcal{G}$ on $\mathcal{A}$ is far from symplectic. But on $\mathcal{A}^\flat$ it acts infinitesimally symplectic. In fact, we have $d_A\xi \in T_A\mathcal{A}^\flat$ for $\xi \in Lie(\mathcal{G})$ and $A \in \mathcal{A}^\flat$, and if we denote by L_ξ the Lie derivative by the fundamental vector field corresponding to $\xi \in Lie\,\mathcal{G}$ we have;

$$\begin{aligned}(L_\xi\omega)_A(a,b) &= (\widetilde{d}\,i_{d_A\xi}\omega)_A(a,b) = \delta_A\,(i_{d_A\xi}\omega_A(b))(a) - \delta_A\,(i_{d_A\xi}\omega_A(a))(b)\\ &= -\frac{1}{24\pi^3}\int_{\partial M} Tr[bd_A\xi - d_A\xi b)a] + \frac{1}{24\pi^3}\int_{\partial M} Tr[ad_A\xi - d_A\xi a)b]\\ &= -\frac{1}{12\pi^3}\int_{\partial M} Tr[(ab-ba)d_A\xi] = -\frac{1}{12\pi^3}\int_{\partial M} d\,Tr[(ab-ba)\xi] = 0,\end{aligned}$$

for $A \in \mathcal{A}^\flat$ and for $a, b \in T_A\mathcal{A}^\flat$.

$\mathcal{G}_0$ being a normal subgroup of $\mathcal{G}$ the action of the quotient group $\mathcal{G}/\mathcal{G}_0$ on $(\mathcal{M}^\flat, \omega^\flat)$ is also infinitesimally symplectic.

Example

The same argument as in Examples 1 and 2 by using paralel transformations along the curves in S^3 yields that

$$\Omega^3 G \simeq \mathcal{A}^\flat(S^3).$$

So we have an injective mapping $\mathcal{M}^\flat(S^3) \longrightarrow \mathcal{A}^\flat(S^3)$. It corresponds to the embedding $\Omega^3_0 G \longrightarrow \Omega^3 G$. Since $\mathcal{M}^\flat(S^3)$ is a fiber bundle over $\mathcal{N}^\flat(S^3)$ with the fiber $\mathcal{G}/\mathcal{G}_0 \simeq \Omega^3_0 G$,

$$\mathcal{N}^\flat(S^3) = \text{ one point.}$$

2 Chern-Simons prequantization over four-manifolds

Let M be an oriented Riemannian four-manifold with boundary and $P = M \times SU(N)$ be the trivial $SU(N)-$ principal bundle. In this section we shall deal with the prequantization of the moduli space of flat connections, that is, over the symplectic manifold $(\mathcal{M}^\flat(\partial M), \omega^\flat)$ we shall construct a line bundle with connection whose curvature is $\omega^\flat$.

2.1 Descent equations

Let $\pi : P \longrightarrow M$ be a principal G-bundle over M. Let Ω^q be the differential q-forms on P and let V^q be the vector space of polynomials $\Phi(A)$ of $A \in \mathcal{A}$

and its curvature F_A that take values in Ω^q. The group of gauge transformations $\mathcal{G}$ acts on V^q by $(g \cdot \Phi)(A) = \Phi(g^{-1} \cdot A)$. We shall investigate the double complex

$$C^{p,q} = C^p(\mathcal{G}, V^{q+3}),$$

that is doubly graded by the chain degree p and the differential form degree q. The coboundary operator $\delta : C^p \longrightarrow C^{p+1}$ is given by

$$\begin{aligned}(\delta\, c^p)(g_1, g_2, \cdots, g_{p+1}) &= g_1 \cdot c^p(g_2, \cdots, g_{p+1}) + (-1)^{p+1} c^p(g_1, g_2, \cdots, g_p) \\ &+ \sum_{k=1}^{p} (-1)^k c^p(g_1, \cdots, g_{k-1}, g_k g_{k+1}, g_{k+2}, \cdots, g_{p+1}).\end{aligned}$$

The following Proposition is a precise version of the Zumino's descent equation. Though stated for $n = 3$ it holds for general n. The first equation is nothing but the Zumino's equation [16]. The author learned the second equation with its transparent proof from Y. Terashima [15]. For $n = 3$ stated here the calculation was appeared in [10, 11].

Proposition 2.1. *Let $c^{0,3} \in C^{0,3}$ be defined by*

$$c^{0,3}(A) = Tr\, F_A^3, \qquad A \in \mathcal{A}.$$

Then there is a sequence of cochains $c^{p,q} \in C^{p,q}$, $0 \le p, q \le 3$ that satisfies the following relations:

$$\begin{aligned} dc^{p,3-p} + (-1)^p \delta c^{p-1,3-p+1} &= 0 \qquad (2.1) \\ dc^{p,2-p} + (-1)^{p-1} \delta c^{p-1,3-p} &= c^{p,3-p} \qquad (2.2) \\ c^{p,q} = 0, \qquad if p + q \neq 2, 3 & \end{aligned}$$

Each term is given by the following form:

$$\begin{aligned} c^{0,2}(A) &= Tr\,(AF^2 - \frac{1}{2}A^3F + \frac{1}{10}A^5). \\ c^{1,2}(g) &= \frac{1}{10} Tr(dg \cdot g^{-1})^5 \\ c^{1,1}(g; A) &= = Tr[-\frac{1}{2}V(AF + FA - A^3) + \frac{1}{4}(VA)^2 + \frac{1}{2}V^3 A] \\ & \qquad \text{where } V = dg\, g^{-1} \\ c^{2,1}(g_1, g_2) &= c^{1,1}(g_2;\, g_1^{-1} dg_1\,) \\ c^{2,0}(g_1, g_2; A) &= -Tr[\frac{1}{2}(dg_2 g_2^{-1})(g_1^{-1}dg_1)(g_1^{-1}Ag_1) - \frac{1}{2}(dg_2 g_2^{-1})(g_1^{-1}Ag_1)(g_1^{-1}dg_1)]. \end{aligned}$$

Remark 2.1. We write here the two-dimensional analogy which is familiar to the readers. In this case we consider the double complex $C^{p,q} = C^p(\mathcal{G}, V^{q+2})$. Put $c^{0,2} = TrF^2 \in C^{0,2}$, then there is a sequence of cochains $c^{p,q} \in C^{p,q}$, $0 \le p, q \le 2$ that satisfies the following relations:

$$\begin{aligned} dc^{0,1} = c^{0,2}, \quad \delta c^{1,0} &= c^{2,0} \\ dc^{1,0} + \delta c^{0,1} &= c^{1,1} \\ dc^{2,0} - \delta c^{1,1} &= 0 \\ c^{p,q} = 0, \qquad \text{if}\, p+q \neq 2, 1 & \end{aligned}$$

Each terms are given as follows.

$$\begin{aligned} c^{0,1}(A) &= Tr\,(AF - \frac{1}{3}A^3). \\ c^{1,0}(A; g) &= Tr(dg\, g^{-1}\, A) \\ c^{1,1}(g) &= \frac{1}{3}Tr(dg\, g^{-1})^3 \\ c^{2,0}(g_1, g_2) &= c^{1,0}(g_1^{-1}dg_1;\, dg_2\, g_2^{-1}) \end{aligned}$$

We note that the relation $dc^{2,0} - \delta c^{1,1} = 0$ represents the Polyakov-Wiegmann formula appeared in [12], and the relation $\delta c^{2,0} = 0$ is the cocycle condition for the central extension of the loopgroup LG.

2.2 Prequantization over $\emptyset = \partial S^4$

2.2.1 4-dimensional Polyakov-Wiegmann formula

Let $G = SU(N)$. We assume in the following that $N \ge 3$. Then $\pi_4(G) = 1$ and $\pi_5(G) \simeq \mathrm{Z}$. Let P be the trivial G-bundle over S^4. Let $\mathcal{A}$ be the space of irreducible connections on P. Let $\mathcal{G}(M)$ be the group of based gauge transformations. We have $\mathcal{G} = S^4G$, the set of smooth mappings from S^4 to G based at some point.

The Chern-Simons form on P is defined by

$$c^{0,2}(A) = Tr\,(AF^2 - \frac{1}{2}A^3F + \frac{1}{10}A^5). \tag{2.3}$$

We have then $Tr(F^3) = d\,c^{0,2}(A)$.

The variation of the Chern-Simons form along the $\mathcal{G}$-orbit is described by the equation (2.2):

$$c^{0,2}(g \cdot A) - c^{0,2}(A) = d\,c^{1,1}(g; A) + c^{1,2}(g). \tag{2.4}$$

Let D^5 be the five dimensional ball with boundary S^4. Then $g \in S^4G$ is extended to D^5G, in fact, we have such an extension by virtue of $\pi_4(G) = 1$.

Put

$$\begin{aligned}
\Gamma(g; A) &= \frac{i}{24\pi^3} \int_{S^4} c^{1,1}(g; A) + C_5(g), \\
C_5(g) &= \frac{i}{24\pi^3} \int_{D^5} c^{1,2}(g) \qquad (2.5) \\
&= \frac{i}{240\pi^3} \int_{D^5} Tr(dg \cdot g^{-1})^5. \qquad (2.6)
\end{aligned}$$

$C_5(g)$ may depend on the extension, but since $\pi_5(G) = \mathrm{Z}$ the difference of two extensions is an integer, and $\exp 2\pi i C_5(g)$ is independent of the extension.

Lemma 2.2 (Polyakov-Wiegmann). *[8, 10, 12] For f, $g \in S^4G$ we have*

$$C_5(fg) = C_5(f) + C_5(g) + \gamma(f, g) \qquad \mathrm{mod}\ \mathbf{Z}, \qquad (2.7)$$

where

$$\begin{aligned}
\gamma(f, g) &= \frac{i}{24\pi^3} \int_{S^4} c^{2,1}(f, g) \\
&= \frac{i}{48\pi^3} \int_{S^4} Tr[(dgg^{-1})(f^{-1}df)^3 + \frac{1}{2}(dgg^{-1}f^{-1}df)^2 + \\
&\qquad + (dgg^{-1})^3(f^{-1}df)].
\end{aligned}$$

proof

From (2,2) we have

$$\delta c^{1,2} + dc^{2,1} = 0.$$

Integrating it over D^5 yields the desired equation. □

Lemma 2.3.

$$\Gamma(fg, A) = \Gamma(g,\ f \cdot A) + \Gamma(f, A) \qquad (2.8)$$

proof

From (2,2) we have

$$\delta c^{0,2} + dc^{1,1} = c^{1,2}.$$

Hence $\delta dc^{1,1} = dc^{2,1}$. Integrating it over D^5, the Polyakov-Wiegmann formula yields the result. □

2.2.2 Prequantum line bundle over $\emptyset = \partial S^4$

We consider the $U(1)-$valued function on $\mathcal{A} \times \mathcal{G}$:

$$\Theta(g, A) = \exp 2\pi i \Gamma(g, A). \tag{2.9}$$

Lemma 2.3 yield the following cocycle condition:

$$\Theta(g, A)\Theta(h, g \cdot A) = \Theta(gh, A). \tag{2.10}$$

Therefore if we define the action of $\mathcal{G}(S^4)$ on $\mathcal{A}(S^4) \times \mathrm{C}$ by

$$(g,\, (A, c)\,) \longrightarrow (g \cdot A,\, \Theta(g, A)c)\,,$$

we have a complex line bundle:

$$\mathcal{L} = \mathcal{A} \times \mathrm{C}/\mathcal{G} \longrightarrow \mathcal{B} = \mathcal{A}/\mathcal{G}. \tag{2.11}$$

Θ being $U(1)$-valued, it is a hermitian line bundle.

We restrict $\mathcal{L}$ to $\mathcal{M}^\flat(\emptyset) = \mathcal{A}^\flat/\mathcal{G}$ and have the line bundle

$$\mathcal{L}(\emptyset) = \mathcal{A}^\flat \times \mathrm{C}/\mathcal{G} \;\longrightarrow\; \mathcal{M}^\flat(\emptyset). \tag{2.12}$$

We call it the *prequantum line bundle* over $\emptyset$; an empty three-manifold.

Since $\mathcal{M}^\flat(\emptyset)$ is one point $\mathcal{L}(\emptyset)$ is isomorphic to the complex line C.

2.2.3 Chern-Simons functional over S^4

Now we introduce the *Chern-Simons functional* CS_{S^4}.

We put, for $A \in \mathcal{A}^\flat(S^4)$,

$$\mathrm{CS}_{S^4}(A) = \frac{i}{24\pi^3}\int_{D^5} c^{0,2}(\widetilde{A}) = \frac{i}{240\pi^3}\int_{D^5} Tr(\widetilde{A}^5), \tag{2.13}$$

where $\widetilde{A}$ is the extension of A to D^5. In general this integral may depend on the extension $\widetilde{A}$, but, as we saw in 1.5, a flat connection A on S^4 is expressed as $A = df\, f^{-1}$ by $f = f_A \in \mathcal{G}(S^4)$, so we have

$$c^{0,2}(A) = \frac{1}{10}Tr(df \cdot f^{-1})^5 = c^{1,2}(f).$$

Hence, if we extend $A = df\, f^{-1}$ to $\widetilde{A} = d\widetilde{f}\,\widetilde{f}^{-1}$ on D^5, we have

$$\mathrm{CS}_{S^4}(A) = \frac{i}{24\pi^3}\int_{D^5} c^{0,2}(\widetilde{A}) = \frac{i}{24\pi^3}\int_{D^5} c^{1,2}(\widetilde{f}) = C_5(f),$$

by virtue of the fact $\pi_5(G) = \mathrm{Z}$ the last quantity is defined independently of the extension of f to D^5 modulo an integer, . Thus $\exp 2\pi i\, \mathrm{CS}_{S^4}(A)$ for $A \in \mathcal{A}^\flat(S^4)$ is well defined.

From (2.7) we have

$$\mathrm{CS}_{S^4}(g \cdot A) = \Gamma(g, A) + \mathrm{CS}_{S^4}(A), \qquad \text{mod. Z},$$

which implies

$$\exp 2\pi i\, \mathrm{CS}_{S^4}(g \cdot A) = \Theta(g, A) \exp 2\pi i\, \mathrm{CS}_{S^4}(A). \tag{2.14}$$

This formula defines a non-vanishing element $[A, \exp 2\pi i\, \mathrm{CS}_{S^4}(A)] \in \mathcal{L}(\emptyset)$. Hence the Chern-Simons functional CS_{S^4} gives the trivialization.

$$\begin{aligned} \mathcal{L}(\emptyset) \quad &\simeq \quad \mathrm{C}, \\ [A, c \exp 2\pi i\, \mathrm{CS}_{S^4}(A)] \quad &\longrightarrow \quad c. \end{aligned} \tag{2.15}$$

Lemma 2.4.

$$\left(\widetilde{d}\, \mathrm{CS}_{S^4}\right)_A (a) = \frac{i}{48\pi^3} \int_{S^4} Tr[A^3\, a], \tag{2.16}$$

for any $A \in \mathcal{A}^\flat(S^4)$.

proof

The lemma follows from the following calculations. For any $A \in \mathcal{A}(S^4)$ and for any extension $\widetilde{A}$ of A to D^5, we have

$$\left(\widetilde{d}\, Tr[\widetilde{A}^5]\right)(a) = 5 Tr[\widetilde{A}^4\, a],$$

and

$$d\, Tr[A^3\, a] = Tr[\, \widetilde{A}^4\, a\,],$$

where $a \in T_{\tilde{A}}\mathcal{A}(D^5)$. And we have

$$\left(\widetilde{d}\, \mathrm{CS}_{S^4}\right)_A (a) = \frac{i}{48\pi^3} \int_{D^5} Tr[\widetilde{A}^4 a] = \frac{i}{48\pi^3} \int_{S^4} Tr[A^3\, a]$$

□

We see from this lemma that if we define the connection form θ on the line bundle $\mathcal{L}(\emptyset)$ by $-2\pi i$ times the right hand side of (2.16) then $\exp 2\pi i\, \mathrm{CS}_{S^4}$ is a horizontal section of the pull-back line bundle $p^*\mathcal{L}(\emptyset)$ by $p : \mathcal{A}^\flat(S^4) \longrightarrow \mathcal{M}^\flat(\emptyset)$.

2.3 Prequantization over 4-dimensional discs

2.3.1 Pre-quantum line bundles over S^3 and $(S^3)'$

Let Ω^3G be the set of smooth mappings from S^3 to $G = SU(N)$ that are based at some point. Ω^3G is not connected but divided into the connected components by deg. We put

$$\Omega^3_0G = \{g \in \Omega^3G;\ \deg g = 0\}. \tag{2.17}$$

The oriented 4-dimensional disc with boundary S^3 is denoted by D, while that with oposite orientation is denoted by D'. The composite cobordism of D and D' becomes S^4. We write $DG = Map(D, G)$, the set of smooth mappings from D to G based at a $p_0 \in \partial D = S^3$, and similarly $D'G = Map(D', G)$. The restriction to S^3 of a $f \in DG$ has degree 0; $f|S^3 \in \Omega^3_0G$. We denote $D_0G = \{f \in DG;\ f|S^3 = 1\}$ and $D'_0G = \{f' \in D'G;\ f'|S^3 = 1\}$. The upper prime will indicate that the function expressed by the letter is defined on D', for example, $1'$ is the constant function $D' \ni x \longrightarrow 1'(x) = 1 \in G$, while 1 is the constant function $D \ni x \longrightarrow 1(x) = 1 \in G$. For $g \in DG$ and $g' \in D'G$ such that $g|S^3 = g'|S^3$, we write by $g \vee g'$ the gauge transformation on S^4 that are obtained by sewing g' and g.

Let P be the trivial $G = SU(N)$-principal bundle over D with $N \geq 3$. For the groups of gauge transformations on D, we have $\mathcal{G}(D) = DG$, $\mathcal{G}_0(D) = D_0G$ and $\mathcal{G}(S^3) = \Omega^3_0G$. Similarly for the trivial G-bundle over D' we have $\mathcal{G}(D') = D'G$ and $\mathcal{G}_0(D') = D'_0G$.

Let $\left(\, \mathcal{M}^\flat(S^3) = \mathcal{A}^\flat(D)/D_0G\, ,\, \omega^\flat \,\right)$ be the moduli space of flat connections over D with the symplectic structure $\omega^\flat$. We shall construct a prequantum line bundle over $\mathcal{M}^\flat(S^3)$. Let $\tilde{P}$ be the trivial extension of P to S^4. We extend any $g \in \mathcal{G}_0(D)$ across the boundary S^3 by defining it to be the identity transformation on $\tilde{P}|D'$. Then $g \vee 1' \in S^4G$. We put, for $A \in \mathcal{A}^\flat(D)$ and $g \in D_0G$,

$$\Gamma_D(g; A) \;=\; \frac{i}{24\pi^3}\int_D c^{1,1}(g; A) + C_5(g \vee 1'),. \tag{2.18}$$

$$\Theta_D(g; A) \;=\; \exp 2\pi i \Gamma_D(g; A). \tag{2.19}$$

$\Theta_D(g, A)$ satisfies the cocycle condition (2.10):

$$\Theta_D(g,\, A)\, \Theta_D(h,\, g\cdot A) = \Theta_D(gh,\, A), \qquad \text{for } g, h \in D_0G.$$

So if we define the action of D_0G on $\mathcal{A}^\flat(D) \times \mathrm{C}$ by

$$(g, (A, c)) \longrightarrow (g\cdot A\, , \Theta_D(g, A)\, c)\, ,$$

we have an hermitian line bundle on $\mathcal{M}^\flat(S^3)$ with the transition function $\Theta(g,A)$:

$$\mathcal{L}(S^3) = \mathcal{A}^\flat(D) \times \mathrm{C}/D_0G \longrightarrow \mathcal{M}^\flat(S^3). \tag{2.20}$$

Now we shall show that the line bundle $\mathcal{L}(S^3)$ has a connection with curvature given by the symplectic form $-i\,\omega^\flat$.

We define the following 1-form θ on $\mathcal{A}^\flat(D)$:

$$\theta_A(a) = \frac{i}{48\pi^3}\int_D Tr[A^3\,a] + \int_D Tr\,\gamma_A^0(a)\,dx, \qquad \text{for } a \in T_A\mathcal{A}^\flat. \tag{2.21}$$

We note that γ^0 is invariant under the action of D_0G. θ is a 0-cochain on the Lie group D_0G taking its value in the space of 1-forms on $\mathcal{A}^\flat(D)$, the coboundary of θ becomes

$$\begin{aligned}(\delta\,\theta_A(a))(g) &= \frac{i}{48\pi^3}\int_D Tr[(g^{-1}Ag + g^{-1}dg)^3 g^{-1}ag - A^3a] \\ &= \widetilde{d}\,\Gamma(g,A)(a) + \frac{i}{48\pi^3}\int_D Tr[(AV^2 + V^2A + 2AVA)a],\end{aligned}$$

for $g \in D_0G$, where $V = dgg^{-1}$. But, since $g \in D_0G$, we have $V|S^3 = 0$, and

$$\int_D Tr[(AV^2 + V^2A + 2AVA)a] = \int_{S^3} Tr[VA - AV)a] = 0,$$

for $a \in T_A\mathcal{A}^\flat(D)$. Therefore

$$(\delta\,\theta_A(a))(g) = \widetilde{d}\,\Gamma_D(g,A)(a) = \frac{1}{2\pi i}(\widetilde{d}\,\log\Theta_D(g,A))(a). \tag{2.22}$$

Thus θ_A gives a connection on the line bundle $\mathcal{L}(S^3)$. The curvature form becomes;

$$\begin{aligned}(\widetilde{d}\,\theta)_A(a,b) &= \frac{i}{24\pi^3}\int_D Tr[A^2(ab - ba)] + \phi_A(a,b) \\ &= \frac{i}{24\pi^3}\int_{S^3} Tr[A(ab - ba)] + \phi_A(a,b) \\ &= -i\,\omega'_A(a,b) + \phi_A(a,b).\end{aligned}$$

From Corollary 1.9 it follows that γ_A^0 and ϕ_A vanishes over $\mathcal{M}^\flat(S^3)$, hence

$$\theta_A(a) = \frac{i}{48\pi^3}\int_D Tr[A^3\,a], \tag{2.23}$$

$$(\widetilde{d}\,\theta)_A(a,b) = -\frac{i}{24\pi^3}\int_{S^3} Tr[A(ab - ba)] = -i\,\omega'_A(a,b), \tag{2.24}$$

for $a,\, b \in T_A\mathcal{M}^\flat(S^3)$.

Thus we have proved the following

Theorem 2.5. *There exists a prequantization of the moduli space* $\big(\mathcal{M}^\flat(S^3),\, \omega^\flat\big)$, *that is, there exists a Hermitian line bundle with connection* $\mathcal{L}(S^3) \longrightarrow \mathcal{M}^\flat(S^3)$, *whose curvature is equal to the symplectic form* $-i\,\omega^\flat$.

Similarly there is a hermitian line bundle with connection $\mathcal{L}((S^3)')$ over the symplectic space $\mathcal{M}^\flat(S^3)$.

$$\mathcal{L}((S^3)') = \mathcal{A}^\flat(D') \times \mathrm{C}/D_0'G \longrightarrow \mathcal{M}^\flat(S^3). \tag{2.25}$$

Remember that we identify $\mathcal{M}^\flat(S^3)$ and $\mathcal{M}^\flat((S^3)')$ by Lemma 1.10. $\mathcal{L}((S^3)')$ is given by the transition function

$$\Theta_{D'}(g', A') = \exp\, 2\pi i \Gamma_{D'}(g', A')$$

for $g' \in D_0'G$ and $A' \in \mathcal{A}^\flat(D')$. Here any $g' \in D_0'G$ is considered to be extended from D' to S^4 by putting it equal to the identity transformation on D.

The connection on $\mathcal{L}((S^3)')$ is given by

$$\theta'_A(a) = \frac{i}{48\pi^3} \int_{D'} Tr[A^3\, a], \qquad \text{for } a \in T_A\mathcal{M}^\flat, \tag{2.26}$$

Then

$$(\widetilde{d}\,\theta')_A(a,b) = -(\widetilde{d}\,\theta)_A(a,b) = i\,\omega_A(a,b).$$

In fact the prequantum line bundles $\mathcal{L}((S^3)')$ and $\mathcal{L}(S^3)$ are in duality.

For $A \in \mathcal{A}(D)$ and $\mathcal{A}(D')$ such that $A|S^3 = A'|(S^3)'$, we write by $A \vee A'$ the connection on S^4 obtained by patching up the two connections along the boundary. We have

$$(g \vee g') \cdot (A \vee A') = (g \cdot A) \vee (g' \cdot A').$$

Now the Polyakov-Wiegmann formula (Lemma 2.2) says

$$C_5(F\,G) = C_5(F) + C_5(G) + \gamma(F, G), \qquad mod\,\mathrm{Z}, \tag{2.27}$$

for F, $G \in \mathcal{G}(S^4)$. Since

$$\gamma(F, G) = \Gamma(F^{-1}dF, G) - C_5(G),$$

we have

$$\Gamma_D(g, A) + \Gamma_{D'}(g', A') = \Gamma(g \vee g', A \vee A'), \qquad mod\,\mathrm{Z},$$

for $g \in D_0G$ and $g' \in D_0'G$. Here we have used the fact $\gamma(g \vee 1', 1 \vee g') = 0$.

Therefore

$$\Theta_D(g,A)\,\Theta_{D'}(g',A') = \Theta_{S^4}(g\vee g', A\vee A'). \tag{2.28}$$

By virtue of this formula we have the homomorphism of line bundles:

$$\mathcal{L}(S^3)\times\mathcal{L}((S^3)') \longrightarrow \mathcal{L}(\emptyset). \tag{2.29}$$

It is given by

$$([A',c'],\,[A,c]) \longrightarrow [A\vee A', c\vee c'], \tag{2.30}$$

where $c\vee c'$ in the right hand side means that $c\in\pi^{-1}(A)$ and $c'\in\pi^{-1}(A')$. Composed with the homomorphism $\mathcal{L}(S^4)\longrightarrow \mathrm{C}$, (2.15), we have the duality of $\mathcal{L}(S^3)$ and $\mathcal{L}((S^3)')$:

$$\mathcal{L}(S^3)\times\mathcal{L}((S^3)') \longrightarrow \mathrm{C}. \tag{2.31}$$

2.3.2 Chern-Simons functional over D and D'

We shall introduce the Chern-Simons functional on D as a section of the pull-back line bundle $p^*\mathcal{L}((S^3)')$ that is horizontal with respect to the connection θ.

We defined in (1.26) the map $p:\mathcal{A}^\flat(D)\longrightarrow\mathcal{M}^\flat((S^3)')$ that induces the identification of $\mathcal{M}^\flat(S^3)$ and $\mathcal{M}^\flat((S^3)')$. Let $A\in\mathcal{A}^\flat(D)$. We consider

$$(A',\,\exp 2\pi i\mathrm{CS}_{S^4}(A\vee A'))\in\mathcal{A}^\flat(D')\times\mathrm{C},$$

for $A'\in\mathcal{A}^\flat(D')$ such that $[A']=p(A)$. Let A'' be another flat connection on D' with $A''|S^3=A|S^3$. There is a $g'\in\mathcal{G}_0(D')$ such that $A''=g'\cdot A'$. So we have $(1\vee g')\cdot(A\vee A')=A\vee A''$. Since

$$\mathrm{CS}_{S^4}(A\vee A'') = \mathrm{CS}_{S^4}(A\vee A') + \Gamma_{D'}(A',g'), \qquad mod.\,\mathrm{Z},$$

it holds that

$$\exp 2\pi i\,\mathrm{CS}_{S^4}(A\vee A'') = \Theta_{D'}(A',g')\cdot\exp 2\pi i\,\mathrm{CS}_{S^4}(A\vee A')\,,$$

so we have a well defined map

$$\exp 2\pi i\,\mathrm{CS}_D\;:\;A\longrightarrow[A',\,\exp 2\pi i\,\mathrm{CS}_{S^4}(A\vee A')]\in\mathcal{L}((S^3)')_{p(A)}, \tag{2.32}$$

that is, we have a section $\exp 2\pi i\,\mathrm{CS}_D$ of $p^*\mathcal{L}((S^3)')$ over $\mathcal{A}^\flat(D)$. We call CS_D the *Chern-Simons functional* over D.

Similarly we have the Chern-Simons functional over D' that is defined by

$$\exp 2\pi i\,\mathrm{CS}_{D'}(A') = [A,\,\exp 2\pi i\,\mathrm{CS}_{S^4}(A\vee A')]\in\mathcal{L}(S^3)_{p'(A')}, \tag{2.33}$$

which is a section of the pull back line bundle $(p')^*\mathcal{L}(S^3)$ by $p' : \mathcal{A}^\flat(D') \longrightarrow \mathcal{M}^\flat(S^3)$.

Let θ be the connection on the line bundle $\mathcal{L}(S^3)$. Let ∇ be the covariant derivative that corresponds to the connection form θ defined by

$$\nabla = \widetilde{d} - i\pi\theta. \tag{2.34}$$

Proposition 2.6. *The functional* $\exp 2\pi i \,\mathrm{CS}_{D'}$ *is a horizontal section of the line bundle* $(p')^*\mathcal{L}(S^3)$*:*

$$\nabla \exp 2\pi i \,\mathrm{CS}_{D'} = 0, \tag{2.35}$$

and it gives the trivialization

$$(p')^*\mathcal{L}(S^3) \simeq \mathcal{A}^\flat(D') \times \mathrm{C}.$$

In fact we have from Lemma 2.2

$$\begin{aligned}(\,\widetilde{d} \exp\, 2\pi i \,\mathrm{CS}_{D'}\,)_A(a) &= \exp 2\pi i \,\mathrm{CS}_{D'}(A)\,(\,\widetilde{d}\,\mathrm{CS}_{D'}\,)_A(a)\\ &= i\pi\theta_A(a)\, \exp 2\pi i \,\mathrm{CS}_{D'}(A).\end{aligned}$$

Similarly, for the pullback line bundle $p^*\mathcal{L}((S^3)') \simeq \mathcal{A}^\flat(D)\times\mathrm{C}$, $\exp 2\pi i \,\mathrm{CS}_D$ is a non-vanishing horizontal section over $\mathcal{A}^\flat(D)$.

2.4 Prequantization over a conformally flat 4-manifold

In the preceeding sections we have defined, for $M = S^4$, D and D',

1. the symplectic structure $\big(\mathcal{M}^\flat(\partial M), \omega^\flat\big)$ on the $\mathcal{G}_0$-orbit space of irreducible flat connections in the trivial bundle $M \times G$.
2. the prequantum line bundle, that is, the hermitian line bundles $\mathcal{L}(\partial M)$ over $\mathcal{M}^\flat(\partial M)$ with connection whose curvature is equal to the symplectic form of $\mathcal{M}^\flat(\partial M)$.
3. the Chern-Simons functional CS_M, such that $\exp 2\pi i\, CS_M$ is a non-vanishing horizontal section of the pull back line bundle of $\mathcal{L}((\partial M)')$ by the boundary restriction map $p : \mathcal{A}^\flat(M) \longrightarrow \mathcal{M}^\flat(\partial M)$. Here the upper prime indicates the opposite orientation.

In the sequel we shall extend functorially these concepts to more general four-manifolds.

First we must make precise what the *Chern-Simons prequantization* means for general four-manifolds. We follow the axiomatic approach proposed for the definition of Topological Field Theory (TFT) by Atiyah [1], and likewise for the Wess-Zumino-Witten model treated in our previous work [8].

Let $\mathcal{X}_4$ be the conformal equivalence classes of all compact conformally flat four-manifolds with boundary.

Let $\mathcal{X}$ be the category whose objects are three-dimensional manifolds Γ which are disjoint union of round S^3's. A morphism between three-dimensional manifolds Γ_1 and Γ_2 is an oriented cobordism given by $\Sigma \in \mathcal{X}_4$ with boundary $\partial\Sigma = \Gamma_2 \bigcup(\Gamma_1')$, where the upper prime indicates the opposite orientation.

Let $\mathcal{Y}$ be the category of complex line bundles.

To each $M \in \mathcal{X}_4$ and $\Gamma = \partial M \in \mathcal{X}$ we have the symplectic space $(\mathcal{M}^\flat(\Gamma), \omega^\flat)$ and the boundary restriction map $p : \mathcal{A}^\flat(M) \longrightarrow \mathcal{M}^\flat(\Gamma) \simeq \mathcal{M}^\flat(\Gamma')$.

A Chern-Simons prequantization means a functor CS from the category $\mathcal{X}$ to the category $\mathcal{Y}$ which assigns:

CS1, to each manifold $\Gamma \in \mathcal{X}$, a complex line bundle $CS(\Gamma)$ over the space $\mathcal{M}^\flat(\Gamma)$,

CS2, to each $M \in \mathcal{X}_4$, with $\partial M = \Gamma$, a section $\exp 2\pi i\, \mathrm{CS}_M$ over $\mathcal{A}^\flat(M)$ of the pullback line bundle $p^*CS(\Gamma)$.

First the functor CS satisfies axioms of TFT. The following axioms **A1**,**A2** and **A3** represent in the category of line bundles the orientation reversal and the operation of disjoint union and contraction [1].

A1 (Involution):

$$CS(\Gamma') = CS(\Gamma)^* \tag{2.36}$$

where $*$ indicates the dual line bundle.

A2 (Multiplicativity):

$$CS(\Gamma_1 \cup \Gamma_2) = CS(\Gamma_1) \otimes CS(\Gamma_2). \tag{2.37}$$

A3 (Associativity):

For a composite cobordism $M = M_1 \cup_{\Gamma_3} M_2$ such that $\partial M_1 = \Gamma_1' \cup \Gamma_3$ and $\partial M_2 = \Gamma_2 \cup \Gamma_3'$, we have

$$\exp 2\pi i\, \mathrm{CS}_M(A) =< \exp 2\pi i\, \mathrm{CS}_{M_1}(A_1), \exp 2\pi i\, \mathrm{CS}_{M_2}(A_2) >, \tag{2.38}$$

for any $A \in \mathcal{A}^\flat(M)$, $A_i = A|M_i$, $i = 1,2$. Where, if we let $p_i : \mathcal{A}^\flat(M_i) \longrightarrow \mathcal{M}^\flat(\Gamma_i) \simeq \mathcal{M}^\flat(\Gamma_i')$, $i = 1,2$, and $p_3^i : \mathcal{A}^\flat(M_i) \longrightarrow \mathcal{M}^\flat(\Gamma_3) \simeq \mathcal{M}^\flat(\Gamma_3')$, $i = 1,2$, then $< , >$ denotes the natural pairing

$$p_1{}^*CS(\Gamma_1') \otimes p_3^{1*}CS(\Gamma_3) \otimes p_3^{2*}CS(\Gamma_3') \otimes p_2{}^*CS(\Gamma_2) \longrightarrow p_1{}^*CS(\Gamma_1') \otimes p_2{}^*CS(\Gamma_2).$$

From **A1** and **A2** it follows that any cobordism M between Γ_1 and Γ_2 induces a homomorphism of sections of pullback line bundles

$$\exp 2\pi i \,\mathrm{CS}_M : C^\infty(p_1^* CS(\Gamma_1)) \longrightarrow C^\infty(p_2^* CS(\Gamma_2)). \tag{2.39}$$

We impose:

$$CS(\phi) = \mathrm{C} \quad \text{for } \emptyset \text{ the empty 3-dimensional manifold.} \tag{2.40}$$

The following axioms characterize the Chern-Simons prequantization.

A4 For each Γ, $CS(\Gamma)$ has a connection with curavature $i\,\omega^\flat$.

A5 $\exp 2\pi i \,\mathrm{CS}_M$ is parallel with respect to the induced connection on $p^* CS(\Gamma)$.

Let $M \in \mathcal{X}_4$ and $\Gamma = \bigcup_{i \in I} \Gamma_i$ be the boundary of M. Each oriented component Γ_i consists of a round S^3, and is endowed with a parametrization $p_i : S^3 \longrightarrow \Gamma_i$. We distinguish positive and negative parametrizations $p_i : S^3 \longrightarrow \Gamma_i$, $i \in I_\pm$, depending on whether p_i respects the orientation of Γ_i or not. We assume further that the compact manifold $\widehat{M}$, that is obtained from M by capping the discs D_i to Γ_i for $i \in I_-$, and D'_i for $i \in I_+$ respectively, is connected and simply connected. $\widehat{M}$ itself is the boundary of a 5 dimensional manifold N; $\partial N = \widehat{M}$, since $\widehat{M}$ is conformally flat.

We define the line bundles $CS(\emptyset)$, $CS(S^3)$ and $CS((S^3)')$ as follows:

$$\begin{aligned} CS(\emptyset) &= \mathcal{L}(\emptyset), \\ CS(S^3) &= \mathcal{L}((S^3)'), \\ CS((S^3)') &= \mathcal{L}(S^3). \end{aligned}$$

The corresponding Chern-Simons functionals are CS_{S^4}, CS_D and $\mathrm{CS}_{D'}$ respectively. They satisfy the axioms **A1**, **A2**, **A4** and **A5**, for $\Gamma = \emptyset$, S^3.

We shall give in the following $CS(\Gamma)$ and CS_M.

$\widehat{M}$ being simply connected $\mathcal{M}^\flat(\widehat{M})$ is one point, which we denote by $\emptyset$. The prequantum line bundle $\mathcal{L}(\emptyset)$ becomes the complex line C. Then the Chern-Simons functional on $\widehat{M}$ is defined as in the case of S^4.

$$\mathrm{CS}_{\hat{M}}(A) = \frac{i}{240\pi^3} \int_N Tr(A^5), \qquad \text{mod.Z.} \tag{2.41}$$

For $i \in I_- \oplus I_+$, the parametrization defines the map $p_i : \mathcal{M}^\flat(\Gamma_i) \longrightarrow \mathcal{M}^\flat(S^3)$. Then we have the pull-back bundle of $\mathcal{L}(S^3)$ (resp. $\mathcal{L}((S^3)')$) by p_i. We define

$$\begin{aligned} \mathcal{L}(\Gamma_i) &= p_i^* \mathcal{L}(S^3) \quad \text{for } i \in I_-, \\ \mathcal{L}(\Gamma_i) &= p_i^* \mathcal{L}((S^3)') \quad \text{for } i \in I_+, \end{aligned} \tag{2.42}$$

then we have respectively

$$\begin{aligned}\mathcal{L}(\Gamma_i') &= p_i^*\mathcal{L}((S^3)') \quad \text{for } i \in I_-,\\ \mathcal{L}(\Gamma_i') &= p_i^*\mathcal{L}(S^3) \quad \text{for } i \in I_+.\end{aligned} \tag{2.43}$$

Here the prime indicates the opposite orientation.

The line bundle $CS(\Gamma)$ is defined by

$$CS(\Gamma) = \otimes_{i\in I_-}\mathcal{L}(\Gamma_i) \otimes \otimes_{i\in I_+}\mathcal{L}(\Gamma_i). \tag{2.44}$$

Now let $\alpha : S^3 \longrightarrow S^3$ be the restriction on S^3 of a conformal diffeomorphism on S^4. First we suppose that α preserves the orientation. Then, since the transition function Θ_D is invariant under α, the line bundle $\mathcal{L}(S^3)$ is invariant under α. If α reverses the orientation then D is mapped to D' and Θ_D is changed to $\Theta_{D'}$. Then $\alpha^*\mathcal{L}(S^3)$ becomes $\mathcal{L}((S^3)')$. On the other hand the parametrizations p_i are uniquely defined up to composition with conformal diffeomorphisms. Therefore $CS(\Gamma)$ is well defined for the conformal equivalence class of $\Gamma \in \mathcal{X}$.

The dual of $CS(\Gamma)$ is

$$CS(\Gamma') = \otimes_{i\in I_-}\mathcal{L}(\Gamma_i') \otimes \otimes_{i\in I_+}\mathcal{L}(\Gamma_i'), \tag{2.45}$$

and the duality; $CS(\Gamma) \times CS(\Gamma') \longrightarrow$ C, is given from (2.30) by:

$$\begin{aligned}&< \otimes_{i\in I_-}[A_i', c_i'] \otimes \otimes_{i\in I_+}[B_i, d_i]\, ,\, \otimes_{i\in I_-}[A_i, c_i] \otimes \otimes_{i\in I_+}[B_i', d_i'] >\\ &= \Pi_{i\in I_-} c_i c_i' \cdot \Pi_{i\in I_+} d_i d_i' \cdot \exp\{-2\pi i \sum_{i\in I_-} \mathrm{CS}_{S^4}(A_i \vee A_i') - 2\pi i \sum_{i\in I_+} \mathrm{CS}_{S^4}(B_i \vee B_i')\}.\end{aligned}$$

$CS(\Gamma)$ is seen to be a Hermitian line bundle with connection. The connection θ is given by the same formula as in (2.23):

$$\theta_A(a) = -\frac{i}{48\pi^3}\int_M Tr[A^3\, a], \qquad \text{for } a \in T_A\mathcal{M}^\flat(\Gamma). \tag{2.46}$$

The curvature of θ is the symplectic form $i\,\omega^\flat$ on $\mathcal{M}^\flat(\Gamma)$.

The Chern-Simons functional over M is defined as a section of the pullback line bundle $p^*CS(\Gamma)$ by the boundary restriction map $p : \mathcal{A}^\flat(M) \longrightarrow \mathcal{M}^\flat(\Gamma)$.

We put

$$\Psi = \otimes_{i\in I_-} \exp\, 2\pi i\, \mathrm{CS}_{D_i} \otimes \otimes_{i\in I_+} \exp\, 2\pi i\, \mathrm{CS}_{D_i'}\, .$$

This is a section of the pullback line bundle of $CS(\Gamma')$ by the boundary restriction map associated to $\partial(\widehat{M}\backslash M) = \Gamma'$, here $\widehat{M}\backslash M = \cup_{i\in I_-} D_i \cup \cup_{i\in I_+} D_i'$.

By the duality between $CS(\Gamma)$ and $CS(\Gamma')$ there exists a section of the pullback line bundle $p^*CS(\Gamma)$. This is by definition the Chern-Simons functional CS_M:

$$\langle\, \Psi(A')\, ,\, \exp\, 2\pi i\, \mathrm{CS}_M(A)\, \rangle = \exp 2\pi i\, \mathrm{CS}_{\hat{M}}(A \vee A'), \tag{2.47}$$

for $A \in \mathcal{A}(M)$ and $A' \in \mathcal{A}(\widehat{M} \setminus M)$.

From the construction the functor CS satisfies the axioms $(A1) - (A5)$.

3 Wess-Zumino-Witten actions on prequantum line bundles

3.1 Four-dimensional WZW model

We saw in 1.6 that the group of gauge transformations $\mathcal{G}(\partial M)$ acts on $\mathcal{M}^\flat(\partial M)$ by an infinitesimal symplectic automorphism. Now we investigate the problem of lifting this action to the line bundle $\mathcal{L}(\partial M)$. For that we consider the Wess-Zumino-Witten line bundles and the associated abelian extesion of the group $\mathcal{G}(\partial M)$ that were introduced in [8, 10]. In 3.1 and 3.2 we shall review briefly the WZW model discussed in our previous work [8].

Let $\mathcal{X}_4$, $\mathcal{X}$ and $\mathcal{Y}$ be as in the previous section. WZW model is a functor WZ from $\mathcal{X}$ to $\mathcal{Y}$ which assigns:

WZ1, to each manifold $\Gamma \in \mathcal{X}$, a complex line bundle $WZ(\Gamma)$ over the space ΓG,

WZ2, to each $\Sigma \in \mathcal{X}_4$, with $\partial\Sigma = \Gamma$, a non-vanishing section WZ_Σ of the pullback line bundle $r^*WZ(\Gamma)$.

Where MG is the set of pointed smooth mappings from a manifold M to G, and $r : \Sigma G \longrightarrow \Gamma G$ is the boundary restriction map.

WZ is imposed to satisfy the TFT axioms,

B1 (Involution):

$$WZ(\Gamma') = WZ(\Gamma)^* \tag{3.1}$$

where $*$ indicates the dual line bundle.

B2 (Multiplicativity):

$$WZ(\Gamma_1 \cup \Gamma_2) = WZ(\Gamma_1) \otimes WZ(\Gamma_2). \tag{3.2}$$

B3 (Associativity): For a composite cobordism $\Sigma = \Sigma_1 \cup_{\Gamma_3} \Sigma_2$ such that $\partial\Sigma_1 = \Gamma_1 \cup \Gamma_3$ and $\partial\Sigma_2 = \Gamma_2 \cup \Gamma_3'$, we have

$$\mathrm{WZ}_\Sigma(f) =< \mathrm{WZ}_{\Sigma_1}(f_1), \mathrm{WZ}_{\Sigma_2}(f_2) >, \tag{3.3}$$

for any $f \in \Sigma G$, $f_i = f|\Sigma_i$, i=1,2.

B1 and **B2** imply that any cobordism Σ between Γ_1 and Γ_2 induces a homomorphism of sections of pullback line bundles

$$WZ(\Sigma) : C^\infty(\Sigma, r_1^* WZ(\Gamma_1)) \longrightarrow C^\infty(\Sigma, r_2^* WZ(\Gamma_2)). \tag{3.4}$$

Besides these TFT axioms we demand that WZ satisfies;

B4 For each Γ, $WZ(\Gamma)$ has a connection.

B5 WZ_Σ is non-vanishing and horizontal with respect to the induced connection on $r^*WZ(\Gamma)$,

B6 For each $\Sigma \in \mathcal{X}_4$ with $\Gamma = \partial\Sigma$, on the pullback line bundle $r^*WZ(\Gamma)$ is defined a multiplication $*$ with respect to which we have

$$\mathrm{WZ}_\Sigma(fg) = \mathrm{WZ}_\Sigma(f) * \mathrm{WZ}_\Sigma(g) \qquad \text{for any } f, g \in \Sigma G. \tag{3.5}$$

We constructed in [8] the WZW model

$$WZ : \mathcal{X} \ni (\Gamma, \Sigma) \Longrightarrow (WZ(\Gamma), \mathrm{WZ}_\Sigma) \in \mathcal{Y}.$$

There the construction of the functor WZ is performed by the same way as we mentioned in 2.6 for the Chern-Simons prequantization functor, that is, after we get ($WZ(S^3)$, WZ_D), ($WZ((S^3)')$, $\mathrm{WZ}_{D'}$) and ($WZ(\emptyset)$, WZ_Σ) for $\Sigma \in \mathcal{X}_4$ such that $\partial\Sigma = \emptyset$, we investigate the relation between the orientation reversion and the duality for the line bundles $WZ(S^3)$, etc., then we have ($WZ(\Gamma)$, WZ_Σ) for $\Gamma \in \mathcal{X}$ so that they satisfy the axioms.

In the following we shall explain after [8] how to construct line bundles $WZ(S^3)$, $WZ(((S^3)')$ and $WZ(\emptyset)$ for $\emptyset = \partial S^4$, and the corresponding WZW functionals; WZ_D, $\mathrm{WZ}_{D'}$ and WZ_{S^4}. We refer [8] for the proof.

3.2 WZW line bundles

We define the action of $\mathcal{G}_0(D) = D_0G$ on $DG \times$ C by

$$h \cdot (f \,, c) = (\, fh \,, \, c\, \Theta_D(h \,, \, f^{-1}df\,)\,, \tag{3.6}$$

for $h \in D_0G$ and $(f, c) \in DG \times \mathrm{C}$.

Remember that, for $A = f^{-1}df \in \mathcal{A}(D)$ and a $h \in D_0G$,

$$\begin{aligned} \Theta_D(h, A) &= \exp 2\pi i\, \Gamma_D(h\,,\, A). \\ \Gamma_D(h\,,\, A) &= \frac{i}{24\pi^3}\int_D c^{1,1}(h,\, A) + C_5(h \vee 1') \\ &= \frac{i}{24\pi^3}\int_D c^{2,1}(f,\, h) + C_5(h \vee 1'). \end{aligned}$$

By definition we put

$$WZ((S^3)') = DG \times \mathrm{C}/D_0G \tag{3.7}$$

It is a complex line bundle over Ω_0^3G. We denote the equivalence class of (f, c) by $[f, c]$, and define the projection

$$\pi : WZ((S^3)') \longrightarrow \Omega_0^3G$$

by $\pi([f, c]) = f|S^3$. The transition function $\chi(f, g)$ for $f, g \in DG$ such that $f|S^3 = g|S^3 \in \Omega_0^3G$ is given by

$$\chi(f, g) = \Theta_D(f^{-1}g\,,\, f^{-1}df). \tag{3.8}$$

We have another bundle on Ω_0^3G . Let

$$WZ(S^3) = D'G \times \mathrm{C}/D'_0G, \tag{3.9}$$

where D'_0G acts on $D'G \times \mathrm{C}$ by

$$h' \cdot (f'\,,\, c') = \left(f'h'\,,\, c'\Theta_{D'}(h',\, f'^{-1}df')\right), \tag{3.10}$$

for $h' \in D'_0G$ and $(f'\,,\, c'\,) \in D'G \times \mathrm{C}$. The projection $\pi : WZ(S^3) \longrightarrow \Omega_0^3G$ is given by $[f', c'] \longrightarrow f'|S^3$. It is a complex line bundle. The transition function $\chi'(f', g')$ for $f', g' \in D'G$ such that $f'|S^3 = g'|S^3$ is given by

$$\chi'(f', g') = \Theta_D(f'^{-1}g\,,\, f'^{-1}df'). \tag{3.11}$$

Where

$$\begin{aligned} \Theta_{D'}(h', A') &= \exp 2\pi i\, \Gamma_{D'}(h',\, A'). \\ \Gamma_{D'}(h',\, A') &= \frac{i}{24\pi^3}\int_{D'} c^{1,1}(h',\, A') + C_5(1 \vee h') \end{aligned}$$

$WZ(S^3)$ and $WZ((S^3)')$ are in duality . To see that, let

$$WZ(\emptyset) = S^4G \times \mathrm{C}/S^4G \tag{3.12}$$

with the action of S^4G on $S^4G\times\mathrm{C}$ defined by $H\cdot(F,c) = (FH,\, c\,\Theta_{S^4}(H,\, F^{-1}dF)\,)$. The duality is given by

$$WZ(S^3)\times WZ((S^3)') \ni (\,[f',c']\,,[f,c]\,) \longrightarrow [\,f\vee f'\,,\, cc'\,\exp 2\pi i\, C_5(f\vee f')\,] \in WZ(\emptyset),$$

composed with the isomorphism similar to (2.15):

$$WZ(\emptyset)\ \ni [\,F, c\exp\{2\pi i C_5(F)\,]\} \longrightarrow\ c\in \mathrm{C}.$$

The WZW action functional WZ_{S^4} is defined by

$$WZ_{S^4}(f) = [f, \exp 2\pi i\, C_5(f)], \qquad \text{for } f\in S^4G. \tag{3.13}$$

Let $r: DG\longrightarrow S^3G$ and $r': D'G\longrightarrow (S^3)'G$ be the boundary restriction maps. The WZW action functional WZ_D is defined as a non-vanishing section of the pullback line bundle $r^*WZ(S^3)$.

$$\mathrm{WZ}_D(f) = [f', \exp 2\pi i\, C_5(f\vee f')\,] \in WZ(S^3)|_{r(f)}, \qquad f\in DG. \tag{3.14}$$

Similarly $\mathrm{WZ}_{D'}$ is defined as a section of $(r')^*WZ((S^3)')$.

$$\mathrm{WZ}_{D'}(f') = [f, \exp 2\pi i\, C_5(f\vee f')\,] \in WZ((S^3)')|_{r'(f')}, \qquad f'\in D'G. \tag{3.15}$$

We have

$$\langle \mathrm{WZ}_D(f),\, \mathrm{WZ}_{D'}(f')\rangle = \mathrm{WZ}_{S^4}(f\vee f'), \tag{3.16}$$

for $f\in DG$, $f'\in D'G$ such that $r(f) = r'(f')$.

The multiplication is defined in $r^*WZ(S^3)$.
For $\lambda = [f', a'] \in WZ(S^3)_{r(f)}$ and $\mu = [g', b'] \in WZ(S^3)_{r(g)}$, we put

$$\begin{aligned}(f,\lambda)\,*\,(g,\mu) &= (f,\,[f',\,a'])\,*\,(g,\,[g',\,b']) \\ &= (fg,\,[\,f'g'\,, a'b'\exp\{2\pi i\gamma(f\vee f', g\vee g')]\,). \end{aligned}\tag{3.17}$$

The right hand side does not depend on the choice of $[f', a']$ and $[g', b']$ but on the equivalence classes λ and μ, and the product is well defined.

Then we have, from Lemma 2.2,

$$\mathrm{WZ}_D(f) * \mathrm{WZ}_D(g) = \mathrm{WZ}_D(fg). \tag{3.18}$$

Similarly we have the product on $(r')^*WZ((S^3)')$ over $D'G$. We omit the product formula which is quite similar to the above.

Now we introduce the action of WZW action functionals on Chern-Simons functionals.

Let $\varphi \in r^*WZ(S^3)$ and $\alpha \in p^*CS(S^3)$. Then $\varphi = (f, [f', a])$ with $f \in DG$, $[f', a] \in WZ(S^3)_{r(f)}$, and $\alpha = (A, [A', c])$ with $A \in \mathcal{A}^\flat(D)$, $[A', c] \in CS(S^3)_{p(A)}$. The action of $r^*WZ(S^3)$ on $p^*CS(S^3)$ is defined by

$$\varphi * \alpha = (\, f \cdot A,\; ac \exp 2\pi i\, \gamma(f \vee f', A \vee A')\,)\,. \tag{3.19}$$

Where, for $F \in S^4G$ and $B \in \mathcal{A}^\flat(S^4)$, we have put

$$\gamma(F, B) = \gamma(F, G) \qquad B = G^{-1}dG.$$

By the same argument as to prove (3.18) using Lemma 2.2 we can prove the following

Theorem 3.1. *1. For $f \in DG$ and $A \in \mathcal{A}^\flat(D)$,*

$$\mathrm{WZ}_D(f) * \exp 2\pi i\, \mathrm{CS}_D(A) = \exp 2\pi i\, \mathrm{CS}_D(f \cdot A). \tag{3.20}$$

2. For $f' \in D'G$ and $A' \in \mathcal{A}^\flat(D')$,

$$\mathrm{WZ}_{D'}(f') * \exp 2\pi i\, \mathrm{CS}_{D'}(A') = \exp 2\pi i\, \mathrm{CS}_{D'}(f' \cdot A'). \tag{3.21}$$

3.3 Abelian extensions of the group $\Omega^3_0 G$

The two-dimensional WZW action gives a central extension $\widehat{LG}$ of the loop group LG. The total space of the $U(1)$-principal bundle $\widehat{LG}$ was described as the set of equivalence classes of pairs $(f, c) \in D^2G \times U(1)$, where D^2 is the 2-dimensional disc with boundary S^1. The equivalence relation was defined on the basis of Polyakov-Wiegmann formula [12], as it was so in our four-dimensional generalization treated above.

However the principal bundle associated to $WZ(S^3)$ or $WZ((S^3)')$ has not any natural group structure contrary to the case of the extension of loop group. Instead J. Mickelsson gave an extension of $\Omega^3_0 G$ by the abelian group $Map(\mathcal{A}_3, U(1))$, where $\mathcal{A}_3$ is the space of connections on S^3 . In the following we shall explain after [8, 10, 11] two extensions of $\Omega^3_0 G$ by the abelian group $Map(\mathcal{A}_3, U(1))$ that are in duality.

We define the action of D_0G on $DG \times Map(\mathcal{A}_3, U(1))$ by

$$h \cdot (f\,, \lambda\,) = (\,fh\,,\, \lambda(\cdot)\Theta_D(h\,,\, f^{-1}df\,)\,, \tag{3.22}$$

for $h \in D_0G$ and $(f,\, \lambda\,) \in DG \times Map(\mathcal{A}_3, U(1))$.

We consider the quotient space by this action;

$$\widehat{\Omega G} = DG \times Map(\mathcal{A}_3, U(1))/D_0G. \tag{3.23}$$

The equivalence class of (f, λ) is denoted by $[f, \lambda]$. The projection $\pi : \widehat{\Omega G} \longrightarrow \Omega_0^3 G$ is defined by $\pi([f\,,\, \lambda]) = f|S^3$. Then $\widehat{\Omega G}$ becomes a principal bundle over $\Omega_0^3 G$ with the structure group $Map(\mathcal{A}_3, U(1))$. The transition function is $\chi(f, g) = \Theta_D(f^{-1}g,\, f^{-1}df)$ for $f,\, g \in DG$ such that $f|S^3 = g|S^3$. Here the $U(1)$ valued function $\chi(\,f\,, g\,)$ is considered as a constant function in $Map(\mathcal{A}_3, U(1))$.

The group structure of $\widehat{\Omega G}$ is given by the Mickelsson's 2-cocycle on D which is defined by the following formula.

$$\begin{aligned} \gamma_D(f, g\,; A) &= \frac{i}{24\pi^3}\int_D (\delta c^{1,1})(\,f,\, g\,;\, A) \\ &= \frac{i}{24\pi^3}\int_{S^3} c^{2,0}(f, g\,;\, A) + \frac{i}{24\pi^3}\int_D c^{2,1}(f, g). \end{aligned} \tag{3.24}$$

We define the multiplication on $DG \times Map(\mathcal{A}_3, U(1))$ by

$$(f,\, \lambda) \bullet (g,\, \mu) = (fg\,,\, \lambda(\cdot)\,\mu_f(\cdot)\exp\, 2\pi i\,\gamma_D(\,f, g;\, A)\,)\,, \tag{3.25}$$

where

$$\mu_f(A) = \mu((f|S^3)^{-1}A(f|S^3) + (f|S^3)^{-1}d(f|S^3)).$$

Then $DG \times Map(\mathcal{A}_3, U(1))$ is endowed with a group structure. The associative law follows from (2.1) and (2.2);

$$\delta\, dc^{2,0} = dc^{3,0} = 0, \qquad \delta\, c^{2,1} = 0.$$

From the definition and Lemma 2.2 we can verify that $[f, \lambda] = [g, \mu]$ if and only if there is a $h \in D_0G$ such that $(g, \mu) = (f, \lambda) \bullet (h, \exp\, 2\pi i\, C_5(h \vee 1'))$, hence the set of elements $(h,\, \exp 2\pi i\, C_5(h \vee 1')\,)$ with $h \in D_0G$ forms a normal subgroup of $DG \times Map(\mathcal{A}_3, U(1))$. Thus $\widehat{\Omega G}$ is endowed with the group structure. Since the group $Map(\mathcal{A}_3, U(1))$ is embedded as a normal subgroup of $\widehat{\Omega G}$, $\widehat{\Omega G}$ is an extension of $\Omega_0^3 G$ by the abelian group $Map(\mathcal{A}_3, U(1))$.

We have another extension of $\Omega_0^3 G$ by $Map(\mathcal{A}_3, U(1))$. We consider

$$\widehat{\Omega' G} = D'G \times Map(\mathcal{A}_3, U(1))/D_0'G, \tag{3.26}$$

where the action of $D_0'G$ is defined by

$$h' \cdot (f', \lambda') = (f'h', \lambda'(\cdot)\Theta_{D'}(h', f'^{-1}df')), \tag{3.27}$$

for $h' \in D_0'G$ and $(f', \lambda') \in D'G \times Map(\mathcal{A}_3, U(1))$. The transition function is $\chi'(f', g')$.

The multiplication on $\widehat{\Omega' G}$ is defined by the same way as above using the Mickelsson's 2-cocycle $\gamma_{D'}(A; f', g')$ on D', and $\widehat{\Omega' G}$ becomes a extension of $\Omega_0^3 G$ by the abelian group $Map(\mathcal{A}_3, U(1))$.

Let the group $Map(\mathcal{A}_3, U(1))$ act on C by $\lambda \cdot c = \lambda(0)c$. Then the associated line bundle to the principal bundle $\widehat{\Omega' G}$ is $WZ(S^3)$, and the line bundle associated to $\widehat{\Omega G}$ is $WZ((S^3)')$.

It is very easy to have a similar abelian extension of the group $\mathcal{G}(\Gamma)$ for $\Gamma \in \mathcal{X}$. In fact $\Gamma \in \mathcal{X}$ being the disjoint union of S^3, we have

$$\mathcal{G}(\Gamma) = \otimes_{I_- \cup I_+} \Omega_0^3 G.$$

Then the extension is given by the product bundle

$$\widehat{\mathcal{G}(\Gamma)} = \otimes_{I_-} \widehat{\Omega G} \otimes \otimes_{I_+} \widehat{\Omega' G} \longrightarrow \mathcal{G}(\Gamma), \tag{3.28}$$

The multiplication (3.25) is extended to $\widehat{\mathcal{G}(\Gamma)}$ by tensor products.

3.4 WZW actions on prequantum line bundles

We shall define the action of $\widehat{\Omega G}$ on $\mathcal{L}(S^3)$.

Let $(f, \lambda) \in DG \times Map(\mathcal{A}_3, U(1))$ and $(A, c) \in \mathcal{A}^\flat(D) \times$ C. We put

$$\beta_D(f, A) = \frac{i}{24\pi^3} \int_D c^{1,1}(f, A). \tag{3.29}$$

Note that the relation $\delta\beta_D = \gamma_D$ holds. If $f \in D_0 G$ then $\Gamma_D(f, A) = \beta_D(f, A) + C_5(f \vee 1')$.

The action of (f, λ) on (A, c) is defined by

$$(f, \lambda) \bullet (A, c) = \left(f \cdot A, c\lambda(A|S^3) \exp 2\pi i\, \beta_D(f, A) \right). \tag{3.30}$$

It is a right action. The relation $\beta_D = \delta\gamma_D$ yields that it defines actually an action:

$$(g, \mu) \bullet ((f, \lambda) \bullet (A, a)) = ((f, \lambda) \bullet (g, \mu)) \bullet (A, a),$$

for (f, λ), $(g, \mu) \in DG \times Map(\mathcal{A}_3, U(1))$ and $(A, a) \in \mathcal{A}^\flat(D) \times$ C.

From the definition of $\mathcal{L}(S^3)$ and the fact that $[h, \exp 2\pi i\, C_5(h \vee 1')]$ for $h \in D_0 G$ gives the unit of $\widehat{\Omega G}$ we see that the above action descends to the action of $\widehat{\Omega' G}$ on $\mathcal{L}(S^3)$. Thus we have proved the following theorem.

Theorem 3.2. *The prequantum line bundle* $\mathcal{L}(S^3)$ *carries an action of* $\widehat{\Omega G}$ *that is equivariant with respect to the infinitesimal symplectic action of* $\Omega_0^3 G$ *on the base space* $\mathcal{M}^\flat(S^3)$.

As for the reduction of $\mathcal{L}(S^3)$ by the action of $\widehat{\Omega G}$ we have the following bundle over $\mathcal{N}^\flat(S^3)$. Let

$$\mathcal{K}(S^3) = \mathcal{A}^\flat(D) \times \mathrm{C}/\widehat{\Omega G},$$

and let $\pi : \mathcal{K}(S^3) \longrightarrow \mathcal{N}^\flat(S^3) = \mathcal{M}^\flat(S^3)/\Omega^3 G$ be the induced projection from $\pi : \mathcal{L}(S^3) \longrightarrow \mathcal{M}^\flat(S^3)$. Then $\pi : \mathcal{K}(S^3) \longrightarrow \mathcal{N}^\flat(S^3)$ becomes a line bundle with the structure group $Map(\mathcal{A}_3, U(1))$. But $\mathcal{N}^\flat(S^3)$ being one point we have

$$\mathcal{K}(S^3) \simeq \mathrm{C}.$$

Similarly there exists an action of $\widehat{\Omega' G}$ on $\mathcal{L}((S^3)')$ that is equivariant with respect to the infinitesimal symplectic action $\Omega_0^3 G$ on $\mathcal{M}^\flat((S^3)') \simeq \mathcal{M}^\flat(S^3)$. In particular $Map(\mathcal{A}_3, U(1))$ acts on $\mathcal{L}((S^3)')$, and the reduction $\mathcal{K}((S^3)') = \mathcal{L}((S^3)')/Map(\mathcal{A}_3, U(1))$ becomes a complex line C.

We can easily generalize the above result to the prequantum line bundle $CS(\Gamma)$ for $\Gamma \in \mathcal{X}$ defined in (2.44). The latter carries the equivariant action of $\widehat{\mathcal{G}(\Gamma)}$ given by the tensor product of the actions (3.30) defined on each factor.

Theorem 3.3. *Let* $\Gamma \in \mathcal{X}$. *The prequantum line bundle* $CS(\Gamma)$ *carries an action of* $\widehat{\mathcal{G}(\Gamma)}$ *that is equivariant with respect to the infinitesimal symplectic action of* $\mathcal{G}(\Gamma)$ *on the base space* $\mathcal{M}^\flat(\Gamma)$. *The reduction of this action becomes*

$$CS(\Gamma)/\widehat{\mathcal{G}(\Gamma)} \simeq \mathrm{C}^d,$$

where d is the number of the boundary components.

References

[1] Atiyah, M. F., *Topological quantum field theories* , Publ. Math. Inst. Hautes Etudes Sci. 68(1989),175-186.

[2] Atiyah, M. F. and Bott, R.,*Yang-Mills equations over Riemann surfaces* , Phil. Trans. R. Soc. Lond. A. 308(1982), 523-615.

[3] Axelrod, S., Della Pietra S. and Witten, E., *Geometric quantization of Chern-Simons gauge theory*, J. Differential Geometry 33 (1991), 787-902.

[4] Bao, D. and Nair, V. P., *A note on the covariant anomaly as an equivariant momentum mapping*, Commun. Math. Phys. (1985), 101, 437-448.

[5] Babelon, O. and Viallet, C. M., *The riemannian geometry of the configuration space of gauge theories*, Commun. Math. Phys. (1981), 81, 515-525.

[6] Donaldson, S. K., *Boundary value problems for Yang-Mills fields, J. Geom. Phys. 8(1992) 89-122.*

[7] Donaldson, S. K. and Kronheimer, P. B. The Geometry of Four-Manifolds Oxford Science Publications, 1990.

[8] Kori, T.,Four-dimensional Wess-Zumino-Witten actions, J. Geometry and Physics 47 (2003), 235-258.

[9] Meinrenken, E. and Woodward, C., Hamiltonian loop group actions and Verlinde factorization, J. Differential Geometry 50 (1998), 417-469.

[10] Mickelson, J. , Current Algebras and Groups, (1989), Plenum Press New York.

[11] Mickelsson, J., Kac-Moody Groups, Topology of the Dirac Determinant Bundle and Fermionization, Commun. Math. Phys. 110(1987),175-183.

[12] Polyakov, A. M. and Wiegmann, P. B., Goldstone fields in two dimensions with multivalued actions, Phys. Lett. B 141, 223-228 (1984).

[13] Ramadas, T. R., Singer, I. M. and Weitsman, J., Some comments on Chern-Simons gauge theory Commun. Math. Phys. 126 (1989), 409-420.

[14] Singer, I. M., The geometry og the orbit space for non-abelian gauge theories, Physica Scripta, Vol. 24(1981), 817-820.

[15] Terashima, Y., Private communication.

[16] Zumino, B., Chiral anomalies and differential geometry, Les Houches Proc.ed. B. S. Dewit and R. Stora (1983),1291-1332.